

ISSN 2224-4980



**International
Journal of Ecosystems and Ecology Science
(IJEES)**

<https://doi.org/10.31407/ijeess>



Essays on Ecosystems and Environmental Research

Volume 12/1, 2022

November 2021 - January 2022

<http://ijeess.net/>

**International
Journal of Ecosystems and Ecology Science
(IJEES)**

<https://doi.org/10.31407/ijeess>



International Journal of Ecosystems and Ecology Science (IJEES), ISSN: 2224-4980

Volume 12, Issue 1, November 2021 - January 2022, (Serial Number 45)



Editor in Chief

Hysen Mankolli, Professor, Ecologist, Plainfield, Illinois, USA; Health and Environment Association, Tirana Albania; Earth System Science Interdisciplinary Center (ESSIC), University of Maryland College Park, USA, Consultant Scientist;

Deputy-Editor-in-Chief

Sukru Dursun, Professor, Konya Technical University, Environmental Engineering Dep., Konya-Turkey;
Massimo Zuchetti, Professor, MIT, Massachusetts Institute of Technology, Cambridge MA, USA;
Cezar Kongoli, Professor, University of Maryland College Park, Earth System Science Interdisciplinary Center (ESSIC), USA;
Selim Dogan, Professor, Konya Technical University, Environmental Engineering Dep., Konya-Turkey;

International Scientific Editorial Board

Sukru Dursun, Professor, Konya Technical University, Environmental Engineering Dep., Konya, Turkey;
Massimo Zuchetti, Professor, MIT, Massachusetts Institute of Technology, Cambridge MA, USA;
Veselin Alexandrov, Professor, National Institute of Meteorology and Hydrology, Bulgaria;
Nicola Senesi, Professor, University of Bari, Bari, Italy;
Wander Michelle M, Professor, University of Illinois at Urbana-Champaign, Urbana, Urbana, IL 61801, USA;
Vladimir Pesic, Professor, University of Montenegro, Department of Biology, Podgorica Montenegro;
Ibraim Dincer, Professor, University of Ontario, Faculty of Engineering and Applied Science, Ontario, Canada;
Hysen Mankolli, Professor, Ecologist, Plainfield, Illinois, USA;
Lyudmyla Symochko, Prof. Assoc. Dr. Uzhhorod National University, Faculty of Biology, Uzhhorod, Ukraine;
Zacharoula S. Andreopoulou, Professor, Aristotle University of Thessaloniki, Thessaloniki, Greece;
Mariana Golumbeanu, PhD, National Institute for Marine Research and Development "Grigore Antipa", Constanta, Romania;
Antonis K. Kokkinakis, Professor, Aristotle University of Thessaloniki, Thessaloniki, Greece;
Olivia Cioboiu, PhD, The Oltenia Museum Craiova, Popa Şapcă str. No 8, Rumania;
Nasser Modirshahla, Professor, Islamic Azad University, Department of Applied Chemistry, Iran;
Nina Liogchii, Professor, Institut of Ecology and Geografy, Republic of Moldova;
Trajce Talevski, Professor, Hydrobiological Institute, Naum Ohidski 50, 6000 Ohrid - R.Macedonia;
Alexander V. Derunkov, Professor, Dep. of Entomology, National Museum of Natural History, Washington DC, USA;
Can Ozgur Colpan, Professor, Ryerson University, Canada;
Muhammad Ashraf, Professor, University of Agriculture, Faculty of Sciences, Faisalabad 38040, Pakistan.;
Roberto Mancinelli, Professor, University of Tuscia, Viterbo, Italy;
Bahrive Gulgun, Professor, Ege University, Izmir, Turkey;
Insaf Mekki, PhD, National Research Institute for Rural Engineering, Water, and Forestry (I.N.R.G.R.E.F), Ariana, Tunisia;
Hany Gaber El Shaer, PhD, IUCN Centre for Mediterranean Cooperation / Global Marine, Egypt;

Ertugrul Esmeray, Ph.D, Karabuk University, Engineering Faculty, Environmental Engineering Department Turkey;
Violeta Vidaček-Hainš, PhD, Ass. Professor, University of Zagreb, Croatia;
Cezar Kongoli, Professor, University of Maryland College Park, Earth System Science Interdisciplinary Center (ESSIC), USA;
Justas Kažys, PhD, Assoc. Professor, Vilnius University, Department of Hydrology and Climatology, Vilnius –Lithuania;
Selim Dogan, PhD, Assoc. Professor Selcuk University, Environmental Engineering Department, Konya Turkey;
Hisham Mostafa Alidrisi, PhD, Assoc. Professor, King Abdulaziz University, Engineering College, Jeddah Saudi, Arabia;
Khalid A. A. Al-Ghamdi, PhD, Assoc. Professor, King Abdulaziz University, Engineering College, Jeddah Saudi, Arabia;
Osman Taylan, PhD, Assoc. Professor, King Abdulaziz University, Engineering College, Jeddah Saudi, Arabia;
Robert J. Kuligowski, PhD, University of Maryland College P, Earth System Science Interdisciplinary Center (ESSIC), USA;
Mariia M. Fedoriak, Professor, Chernivtsi National University, Dep. of Ecology and Biomonitoring, Chernivtsi, Ukraine;
Narayan Ramappa Birasal, Associate Professor, KLE Society's G H College, Zoology Department, , Karnataka state, India;
Rana P. Singh, Professor, Babasaheb Bhimrao Ambedkar University, Dep. of Environmental Science, U.P., India;
Hayati AKMAN, Dr., Selcuk University, Seed Department, 42430 Konya, Turkey;
Fatma ÇANKA KILIÇ, Professor, Kocaeli University, Technology Faculty, Energy Systems Engineering Dep., Kocaeli, Turkey;
Ferim Gashi, Dr., University of Pristina, Department of Geography, Faculty of Mathematical-Natural Sciences, Kosovo;
Altin Dorri, Professor Assoc., Polytechnic University of Tirana, Tirana, Albania;
Alexander P. Sizykh, Professor, Russian Academy of Sciences, Siberian Institute, Irkutsk, Lermonyova, Russia;
Zeynep EREN, Professor, Ataturk University, Environmental Engineering Department, Erzurum, Turkey;
Sheida Korjani, PhD, Department of Architecture, Tabriz Branch, Islamic Azad University, Tabriz, Iran;
Maan Maarroof, PhD, Assistant Professor, University of Mosul, College of Veterinary Medicine, Iraq;
Koula Doukani, Professor, University of Ibn Khaldoun, Zaaroura, Tiaret, Algeria;
Svetla Petkova Gateva, PhD, Institute of Biodiversity and Ecosystem Research, Bulgarian Acad. of Sciences, Sofia, Bulgaria;
Rrahim Sejdiu, Professor Assistant, PhD, University of Applied Sciences-Ferizaj, Kosovo;
Mouloud BOUHOUBOU, Dr., ENS Assia Djebar - Constantine, Algeria;
Dinh Tran Ngoc Huy, Dr., Banking University HCMC, Ho Chi Minh city Vietnam-International University of Japan, Japan;
Nataliia P. Kovalenko, PhD, Inst. of History of Agrarian Science, Education and Technique of NSAL NAAS, Kiev, Ukraine;

Publication Information:

International journal of ecosystems and ecology science (IJEES), ISSN 2224-4980, publishing original articles, reviews and short communications of high scientific standard on ecosystems and science ecology. By: Hysen Mankolli, Professor, Ecologist, Plainfield, Illinois, USA; Health and Environment Association, Tirana, Albania;

ISSN International Centre, *Bibliographic Data Section*, PARIS, FRANCE;

Bib - ID 68337; ISO standard 3297;

Key title: International journal of ecosystems and ecology science

Abbreviated key title: *Int. j. ecosyst. ecol. sci.*

ISSN 2224-4980; <http://ijeess.net/>

Frequency: four times a year. IJEES Electronic Journal Publication: Plainfield, Illinois, USA;

DOI prefix: 10.31407, <https://doi.org/10.31407/ijeess>



ESCI Clarivate, [http://ip-science.thomsonreuters.com/cgi-](http://ip-science.thomsonreuters.com/cgi-bin/jrnlst/jlresults.cgi?PC=MASTER&Word=ijeess)

[bin/jrnlst/jlresults.cgi?PC=MASTER&Word=ijeess](http://ip-science.thomsonreuters.com/cgi-bin/jrnlst/jlresults.cgi?PC=MASTER&Word=ijeess)

UOI license: <http://u-o-i.org/1.01/ijeess>; <http://www.u-o-i.org/index.php/SearchLicense>



[DPI Digital Library](http://www.dpi.com) USA and assign unique digital no. DPI: 16.10047.IJEES

U.S. National Library of Medicine,

NLM ID: [101726509](https://www.ncbi.nlm.nih.gov/nlmcatalog/101726509) [Serial]; **Other ID:** (OCOLC)879805056;

<https://www.ncbi.nlm.nih.gov/nlmcatalog/101726509>

This journal has the status of an international journal.

Aims and Scope:

The goals of the International Journal of Ecosystems and Ecology Science (IJEES), ISSN 2224-4980, are to bring together researchers and scientists with interests in the quality of the ecosystems research results, theories, technologies, systems, tools, applications, the work in progress and experiences on ecosystems used. The main topics of interest are:

- ✓ Ecosystems
- ✓ Agro ecosystems
- ✓ Forest ecosystems
- ✓ Ecology
- ✓ Plant Ecology
- ✓ Animal Ecology
- ✓ Human Ecology
- ✓ Lakes and Rivers ecosystems
- ✓ Applied Biology
- ✓ Applied Ecology
- ✓ Applied Chemistry
- ✓ Biodiversity
- ✓ Energy
- ✓ Geology
- ✓ Bioaccumulation
- ✓ Desalination
- ✓ Water
- ✓ Soil
- ✓ Air pollution
- ✓ Climate Change
- ✓ Ecosystem restoration
- ✓ Environment toxicology
- ✓ Environment protection
- ✓ Environmental radioactivity
- ✓ Environmental legislation
- ✓ Environmental management
- ✓ Environmental education.
- ✓ Green technology
- ✓ Socioeconomic aspects in Ecosystems
- ✓ Agro tourism and National Park
- ✓ Health Care
- ✓ Food security
- ✓ Rights and Environmental Laws
- ✓ Environment Engineering
- ✓ Environment Architecture
- ✓ Eco philosophy Issue

Abstracted / Indexed / Impact Factor in:

International journal of ecosystems and ecology science (IJEES), ISSN 2224-4980,

IJEES: <http://ijeess.net/>

IJEES have an international status and has been indexed, abstracted, and have impact factors:

ISSN Center responsible of the record: CIEPS - ISSN

<https://portal.issn.org/resource/ISSN/2224-4980>

IJEES doi crossref

<https://doi.org/10.31407/ijeess>

IJEES, Volume 12/1, 2022

[https://www.ijeess.net/journal-83-International--Journal-of-Ecosystems-and-Ecology-Science--\(IJEES\)--Volume-12-1,-2022.html](https://www.ijeess.net/journal-83-International--Journal-of-Ecosystems-and-Ecology-Science--(IJEES)--Volume-12-1,-2022.html)

Clarivate Analytics

<http://ip-science.thomsonreuters.com/cgi-bin/jrnlst/jlresults.cgi?PC=MASTER&Word=ijeess>

<https://mjl.clarivate.com/home?PC=MASTER&Word=ijeess>

Web of Science Core Collection

<https://apps.clarivate.com/mjl-beta/search-results>

Emerging Sources Citation Index - JOURNAL LIST

<http://mjl.clarivate.com/cgi-bin/jrnlst/jlresults.cgi?PC=EX&Alpha=I>

Sciencegate cited

<https://www.sciencegate.app/source/315335/top-cited>

Scilit

<https://www.scilit.net/journal-articles/1892436>

CiteFactor: IF 2019-2020:1.40;

<https://www.citefactor.org/impact-factor/impact-factor-of-journal-International-Journal-of-Ecosystems-and-Ecology-Science-IJEES.php>

EBSCO's library products,USA

<http://www.ebscohost.com/for-publishers/for-publishers>

IARC: Journal Impact Factor (USA), IF=2017: 1.811; IF=2016: 1.463; =2015: 0.861

http://iarcif.org/index.php/journal_information/show/44/0/2224-4980

Index Copernicus ICV= 2020: 100.00; ICV= 2019: 100.00; ICV= 2018: 100.00; ICV= 2017: 98.95;

<https://journals.indexcopernicus.com/search/journal/issue?issueId=197247&journalId=31407>

Academic Resource Index Research Bible:

<http://paper.researchbib.com/view/issn/2224-4980/10/2>

Google scholar Cited

<https://scholar.google.com/citations?user=W89N8qgAAAAJ&hl=en>

MIAR

<http://miar.ub.edu/issn/2224-4980>

Directory of Research Journals Indexing (DRJI)

<http://olddrji.lbp.world/JournalProfile.aspx?jid=2224-4980>

WorldCat

https://www.worldcat.org/search?qt=affiliate_wc_org_all&ai=Directory_ashok.drji%252540gmail.com&fq=&q=2224-4980&wcsbtn2w=Go

US. National Library of Medicine, NLM ID: 101726509 [Serial]; Other ID: (OCoLC)879805056;

<https://www.ncbi.nlm.nih.gov/nlmcatalog/101726509>

Global Impact Factor (GIF) IF= 2015: 0.765; 2014: 0.625; 2013: 0.541; IF= 2012: 0.425;

<http://globalimpactfactor.com/journals-list/?snap=I>

CAS Chemical Abstracts Service – CAS database

<http://cassi.cas.org/search.jsp>

Journal Factor: ISSN 2224-4980

http://www.journalfactor.org/advance_search.php

Directory of Research Journals Indexing (DRJI)

<http://olddrji.lbp.world/justIncluded.aspx>

Environment-and-Nature/Biodiversity/Ecosystem.

<http://www.efita.org/Environment-and-Nature/Biodiversity/Ecosystem-Biodiversity/IJEES-details-28892.html>

NewJour : Electronic Journal&Newsletters

<http://gulib.georgetown.edu/newjour/nj2/index.html>

Environmental expert

<http://www.environmental-expert.com/magazines/?keyword=ecosystem+science>

Currently Indexing at Pro Quest

http://www.proquest.com/en-US/catalogs/databases/detail/periodicals_index.shtml

CABI

<http://www.cabi.org/default.aspx?page=1016&site=170&pid=40&xslttab=2&newtitlesonly=1&search=>

Manuscripts and correspondence are invited for publication

*Dear Professors, PhD, researchers and scholars of the field of ecosystems and related areas to, such as environmental engineering, agriculture, bio-chemistry, soil, water and air pollution, geology, forestry, urban ecology, resource economics, etc., at the moment, I declare the sending of scientific journal articles to International Journal of Ecosystems and Ecology Science (IJEES) is open. Sending of items can only be done electronically, to the following address: e-mail: editorijeess@gmail.com; 2011ijeese@gmail.com; mjedisishendeti@yahoo.com; The confirmation of receiving these items will be announced within the period of one to three weeks after sending. Whether the article is accepted or not, will be announced within a period of two to three weeks. Only original articles will be accepted, validated, and not to have or be published in other journals. Publication lasts from two to six months. All Rights Reserved Copyright by © IJEES (the first or correspondent author have responsible for the accuracy of the article). Submission guidelines and Web Submission System are available at: <https://www.ijeess.net/page-6-Guide-for-authors.html>;
Note: If the paper is published in another journal or if it includes false or inaccurate claims, it will be rejected or removed automatically as that constitutes a violation. Zero tolerance!*

Editorial Office:

*IJEES Electronic Journal Publication: Plainfield, Illinois, USA; E-mail: editorijeess@gmail.com;
Health and Environment Association, Tirana, Albania; E-mail: mjedisishendeti@yahoo.com;*

**International
Journal of Ecosystems and Ecology Science
(IJEES)**

DOI: <https://doi.org/10.31407/ijeess>

Volume 12/1, 2022

DOI: <https://doi.org/10.31407/ijeess12.1>

Table of contents:

Alexander Vasilyevich Baranovsky¹, Alexey Sergeevich Sadovoy¹, Sergey Ivanovich Kapustin^{2,3}, Andrey Sergeevich Kapustin^{4*}, Alexander Mikhailovich Stroyny¹, Anna Sergeevna Golub³, CHARACTERISTICS OF CONSUMPTIVE WATER USE OF MILLET AND SORGHUM DEPENDING ON THE SOWING TIME IN DRY CONDITIONS OF STEPPE ZONE, page 1-6;

DOI: <https://doi.org/10.31407/ijeess12.101>

Fatima-Zohra Kenza Labbani^{1,2,*}, Scheherazad Dakhmouche^{1,2}, Leila Bennamoun^{2,3}, Amel Ait-Kaki^{2,4}, Tahar Nouadri^{2,3}, EXTRACELLULAR HYDROLYTIC ENZYMES OF YEASTS ISOLATED FROM FRUIT AND BEET PEELS IN ALGERIA, page 7-16;

DOI: <https://doi.org/10.31407/ijeess12.102>

Julinda Jaho^{1*}, Suzana Nuellari², Artan Bano³, DECOMPENSATION OF CHRONIC OPEN ANGLE GLAUCOMA FOLLOWING A SPINAL STEROID INJECTION: A CASE REPORT, page 17-20;

DOI: <https://doi.org/10.31407/ijeess12.103>

Arpit Sharma^{1*}, Sanjeevi Shanthakumar², ACCOUNTABILITY OF CORPORATE TOWARDS ENVIRONMENTAL ISSUES THROUGH THE LENS OF CORPORATE SOCIAL RESPONSIBILITY (FINANCIAL) AND BUSINESS RESPONSIBILITY (NON-FINANCIAL) REGULATIONS WITH REFERENCE TO TOP 30 COMPANIES ON NIFTY, page 21-28;

DOI: <https://doi.org/10.31407/ijeess12.104>

Arjan Korpa^{1*}, Sara Dervishi¹, Diana Gecaj¹, Kristi Shahu¹, Spiro Drushku², REGENERATION EFFICIENCY ASSESSMENT OF A UMO TREATED WITH ACTIVATED KARAEVA'S KOSOVO BENTONITE, page 29-40;

DOI: <https://doi.org/10.31407/ijeess12.105>

Viacheslav Sichkar¹, Volodymyr Orekhivskyi², Lyudmila Bilyavskaya³, Anna Kryvenko^{1*}, Ruslan Solomonov¹, Anna Diyanova³, USE OF SOYBEAN GENETIC RESOURCES TO CREATE HIGHLY ADAPTIVE VARIETIES, page 41-58;

DOI: <https://doi.org/10.31407/ijeess12.106>

Lulzime Dhora^{1*}, THE CONDITIONS OF MARKET FISH POPULATIONS, THAT IMPACT ON SHKODRA LAKE STABILITY, WATER QUALITY AND SUSTAINABLE FISHING, page 59-66;

DOI: <https://doi.org/10.31407/ijeess12.107>

Risida Gjonej^{1*}, Albana Poloska¹, Etleva Smakaj¹, Valbona Bezhani¹, Flora Zyberaj¹, ASSOCIATION OF DELIVERY MODE AND BREASTFEEDING, page 67-76;

DOI: <https://doi.org/10.31407/ijeess12.108>

Heyam Abdulrahman Al Mousa¹, Haseebullah Abdul Naeem^{1*}, Zainab Mohammed Alwan Al- Juboori², PERCEIVED-HOTELSCAPES AND ITS IMPACT ON CUSTOMER PATRONAGE: AN EMPIRICAL PERSPECTIVES, page 77-86;

DOI: <https://doi.org/10.31407/ijeess12.109>

Nataliia Kovalenko^{1*}, Svitlana Yehorova², THE HISTORY OF DEVELOPMENT AND THE PERSPECTIVES OF IMPLEMENTATION OF INNOVATIVE TECHNOLOGIES FOR THE ENVIRONMENTALLY SAFE USE OF WATER AND LAND RESOURCES IN UKRAINE AT THE BEGINNING OF THE 21st CENTURY, page 87-100;

DOI: <https://doi.org/10.31407/ijeess12.110>

Muhamet Kadrija^{1*}, Fatmira Karabollaj², LEVEL OF KNOWLEDGE AND NURSING PRACTICES ON DIABETIC FOOT MANAGEMENT, page 101-106;

DOI: <https://doi.org/10.31407/ijeess12.111>

Albert Lama¹, Diamant Shtiza^{2*}, Enkelejda Shkurti³, THE IMPORTANCE OF URINE CULTURES IN THE FOLLOW-UP AFTER FIRST FEBRILE UTI, page 107-110;

DOI: <https://doi.org/10.31407/ijeess12.112>

Nguyen Thi Hoa^{1*}, Dinh Tran Ngoc Huy², Tran Van Trung¹, IMPLEMENT POLICY ON STUDENT SCIENTIFIC RESEARCH AT VIETNAM UNIVERSITIES AND EDUCATIONAL INSTITUTIONS TODAY: ISSUES TO DISCUSS, page 111-118;

DOI: <https://doi.org/10.31407/ijeess12.113>

Safwan Al Salaimh^{1*}, Khaldoun Besoul², Ayman Nayef Al Halaybeh², POWERFUL ALGORITHM FOR ADAPTIVE RECOGNITION OF DYNAMIC SYSTEM PARAMETERS, page 119-124;

DOI: <https://doi.org/10.31407/ijeess12.114>

Andia Meksi¹, Enkelejda Shkurti^{2*}, ORGANIZATIONAL, ETHICAL AND LEGAL ASPECTS OF PALLIATIVE CARE IN ALBANIA, page 125-132;

DOI: <https://doi.org/10.31407/ijeess12.115>

Ekaterina Alexandrovna Vetrova^{1*}, Elena Evgen'evna Kabanova², Elena Vladimirovna Dupliy³, Julia Olegovna Sulyagina³, Sergey Anatolyevich Makushkin³, THE LANDSCAPING FEATURES OF THE CITY OF BALASHOV, RUSSIA, page 133-140;

DOI: <https://doi.org/10.31407/ijeess12.116>

Adam Abdeljalil^{1*}, Saffaj Nabil¹, Mamouni Rachid¹, TOWARDS A GUIDELINE OF A SPILL MANAGEMENT: INDUSTRIAL SITES AS A CASE STUDY, page 141-148;

DOI: <https://doi.org/10.31407/ijeess12.117>

Sergushina Elena Sergeevna^{1*}, Kabanov Oleg Vladimirovich¹, Lazareva Alina Igorevna¹, Aimen Anuarbek Talhauy², Atasheva Darya², Lyalin Evgeny Alexandrochich³, Pichugin Dmitry Vladimirovich⁴, Baulin Mikhail Aleksandrovich⁴, Bartsaykin Alexey Viktorovich⁵, MARKET ASSESSMENT OF PAYMENT FOR ECOSYSTEM SERVICES, page 149-158;

DOI: <https://doi.org/10.31407/ijeess12.118>

Yana A. Volynchuk^{1*}, Tatiana D. Lykova², Vasilina A. Popil², POSSIBILITIES OF TERRITORIAL BRANDING AS DEVELOPMENT MANAGEMENT TOOL FOR THE FAR EAST OF RUSSIA, page 159-164;

DOI: <https://doi.org/10.31407/ijeess12.119>

Valentina V. Zhokhova^{1*}, Natalya P. Belozertseva¹, Natalya A. Yurchenko¹, FORECAST ESTIMATES OF FAR EASTERN EXPORT TREPANG TO CHINA, page 165-178;

DOI: <https://doi.org/10.31407/ijeess12.120>

Alla A. Udovikova^{1*}, Irina N. Marchenkova¹, A PRACTICE-ORIENTED APPROACH TO THE FINANCIAL STATEMENT-BASED BANKRUPTCY DIAGNOSIS, page 179-186;
DOI: <https://doi.org/10.31407/ijeess12.121>

Larisa A. Glazunova^{1*}, Angelina A. Yurchenko¹, Yuri V. Glazunov¹, Evgenii M. Gagarin¹, Ivan V. Plotnikov¹, CONTAMINATION OF RAW MILK WITH CONDITIONALLY PATHOGENIC MICROORGANISMS AND ANTIBIOTICS, page 187-194;
DOI: <https://doi.org/10.31407/ijeess12.122>

D.Iu. Tiulin^{1*}, A.A. Vasiliev¹, Iu.A. Guseva¹, O.A. Gurkin¹, A.A. Anurieva¹, FISH REPRODUCTION CONDITIONS OF THE VOLGOGRAD RESERVOIR IN THE WATER AREAS NEAR THE VILLAGES OF AKHMAT AND ZOLOTOTIE IN 2020 IN COMPARISON WITH PREVIOUS YEARS, page 195-200;
DOI: <https://doi.org/10.31407/ijeess12.123>

Mikhail Yu. Tretiakov^{1*}, Valeriy K. Tokhtar¹, Anna E. Ivleva¹, Julia N. Kurkina¹, Irina V. Batlutskaya¹, Igor V. Lyashenko¹, OPTIMIZING THE DRY PLANT SAMPLE PREPARATION METHOD FOR EFFICIENT DNA EXTRACTION IN *SYRINGA VULGARIS*, page 201-206;
DOI: <https://doi.org/10.31407/ijeess12.124>

Valeriy K. Tokhtar^{*}, Mikhail Yu. Tretiakov, Victoria N. Zelenkova, Tatiana V. Petrunova, ASSESSMENT OF THE PHYTOREMEDIATION POTENTIAL OF AQUATIC PLANTS OF THE BELGOROD REGION FOR WASTEWATER TREATMENT, page 207-216;
DOI: <https://doi.org/10.31407/ijeess12.125>

Elena Igorevna Artemova¹, Olga Yuryevna Voronkova², Alexey Vladimirovich Kulchitsky³, Tatiana Victorovna Morozova⁴, Ravil Gabdullaevich Akhmadeev^{4*}, Vladimir Mikhailovich Plosky⁵, COUNTRY ASPECTS OF USE OF TAX INSTRUMENTS ON THE HYDROCARBON MARKET, page 217-224;
DOI: <https://doi.org/10.31407/ijeess12.126>

Behcet Oznacar^{1*}, Mahmut Celik¹, FACTORS THAT LEAD TO CHANGE IN THE MISSION AND VISION STATEMENTS OF ORGANIZATIONS: A RESEARCH ON FIVE-STAR HOTELS OPERATING IN THE TURKISH REPUBLIC OF NORTHERN CYPRUS, page 225-236;
DOI: <https://doi.org/10.31407/ijeess12.127>

Mustafa GURSOY^{1*}, BUSINESS ORGANIZATIONAL RESPONSE TO ENVIRONMENTAL ISSUES, page 237-244;
DOI: <https://doi.org/10.31407/ijeess12.128>

Alexander M. Konstantinov^{1*}, EXPLORING THE ESSENCE OF THE GAME IN CONTEXT INTERACTION OF GAMING AND LEGAL NORMS, page 245-252;
DOI: <https://doi.org/10.31407/ijeess12.129>

Ivan A. Usenkov^{1*}, EVALUATION OF “SUNSET LEGISLATION” AS A MEANS OF DEREGULATION, page 253-260;
DOI: <https://doi.org/10.31407/ijeess12.130>

Galina Vukovich^{1*}, Anastasiya Ostrovskaya¹, Lina Zakharova¹, Egor Ponomarenko¹, Ekaterina Slepceva¹, STRATEGIC TRENDS THAT DETERMINE THE ACTUALIZATION OF MARKETING ELEMENTS OF PERSONNEL IN THE FIELD OF AGROTOURISM IN RUSSIA DURING THE CORONACRISIS, page 261-272;
DOI: <https://doi.org/10.31407/ijeess12.131>

Galina Vukovich^{1*}, Anastasiya Ostrovskaya¹, Lina Zaharova¹, Albert Kovalenko¹, Valentina Kuznecova², FORMALIZATION OF THE MARKETING CONTOUR HR-POLICIES IN THE FIELD OF AGROTOURISM IN RUSSIA AS A CONDITION FOR ITS BALANCED DEVELOPMENT, page 273-282;
DOI: <https://doi.org/10.31407/ijeess12.132>

Krystel Grace V. Padilla^{1*}, SCREENING OF PLANT-DERIVED EXTRACTS AGAINST DNase PRODUCTION OF *Staphylococcus aureus* 1582, page 283-290;
DOI: <https://doi.org/10.31407/ijees12.133>

Razanov Serhii^{1*}, Landin Volodymyr², Nedashkivskyi Volodymyr³, Ohorodnichuk Halina⁴, Gucol Galina⁵, Symochko Lyudmyla⁶, Komynar Mykola⁷, INTENSITY OF ¹³⁷CS TRANSITION INTO NECTAR-POLLINATING PLANTS AND BEEKEEPING PRODUCTS DURING RECLAMATION OF RADIOACTIVELY CONTAMINATED SOILS, page 291-298;
DOI: <https://doi.org/10.31407/ijees12.134>

Danny O. Alfonso^{1*}, MYCOCHEMICAL SCREENING, PROXIMATE ANALYSIS AND ANTIMICROBIAL ACTIVITY OF *Pleurotus pulmonarius* (Fr.) Quel, page 299-304;
DOI: <https://doi.org/10.31407/ijees12.135>

Olga Hafiak¹, Lyudmyla Symochko^{1,2*}, SOIL AND WATER MICROBIOTA AS BIOINDICATORS FOR THE ASSESSMENT ECOLOGICAL STATUS OF ECOSYSTEMS, page 305-312;
DOI: <https://doi.org/10.31407/ijees12.136>

INTENSITY OF ^{137}Cs TRANSITION INTO NECTAR-POLLINATING PLANTS AND BEEKEEPING PRODUCTS DURING RECLAMATION OF RADIOACTIVELY CONTAMINATED SOILS

Razanov Serhii^{1*}, Landin Volodymyr², Nedashkivskiy Volodymyr³, Ohorodnichuk Halina⁴,
Gucol Galina⁵, Symochko Lyudmyla⁶, Komynar Mykola⁷

^{1*,4,5}Vinnitsia National Agrarian University, 3, Soniachna Str., Vinnitsia, Ukraine;

²Institute for Safety Problems of Nuclear Power Plants of the National Academy of Sciences of Ukraine,
12, Lisogorskaya Str., Kiev, Ukraine;

³Bila Tserkva National Agrarian University, 8/1, Soborna pl., Bila Tserkva, Ukraine;

⁶Uzhhorod National University, Voloshyna Str.3., Uzhhorod, Ukraine;

⁷Institute of Agroecology and Environmental Management NAAS, 12, Metrologicheskaya Str., Kiev, Ukraine;

*Corresponding Author Razanov Serhii, email: razanovsergej65@gmail.com;

Received November 2021; Accepted December 2021; Published January 2022;

DOI: <https://doi.org/10.31407/ijeess12.134>

ABSTRACT

The article presents investigations of transition ^{137}Cs to agricultural crops (sunflower) and beekeeping products (honey, bee pollen) for reclamation of radioactively contaminated soils. It is proved that the quality of beekeeping products depends from the ecological condition of nectar-pollinating lands. As a result of the 1986 accident at the Chernobyl nuclear power plant, nectar-pollinating lands were subjected to high man-caused impact, in particular, some areas of Polissya. It caused a certain accumulation of ^{137}Cs and ^{90}Sr in beekeeping products. The authors studied the effect of reclamation of contaminated soils, in particular, in the Narodyt'skyi district of Zhytomyr region of Ukraine with ^{137}Cs up to 5 Ci/m^2 with different acidity on the intensity of accumulation of this radionuclide in honey and bee pollen produced by bees from nectar and sunflower pollen. It was found that the specific activity of ^{137}Cs in honey and bee pollen produced by bees from nectar and pollen of sunflower grown on agricultural land with a content of this radionuclide in soils from 2824 Bq/kg to 2665 Bq/kg, not exceeding DR-2006 200 Bq/kg. Reclamation of radioactively contaminated soils with hydrolytic acidity from 1.6 mg to 2.4 mg-eq/100 g of soil, in particular, the application of defecation mud in them at a rate of 4 t/ha to 6 t/ha reduced the specific activity and accumulation coefficient of ^{137}Cs in vegetative mass of sunflower, honey and bee pollen, made by bees from nectar and pollen of this plant.

Keywords: sunflower, soil, honey, bee pollination, vegetative mass, nectar, pollen, specific activity of ^{137}Cs .

INTRODUCTION

Nectar-pollinating plants play an important role in the existence of the honey bee. Nectar and pollen are a source of food for bees and raw materials for their products, including honey, perga, wax, royal jelly, homogenate of drone larvae and others (Nicolson, 2011).

The nectar-pollinating base of Ukraine includes about 900 species of plants. Among this number of nectar pollinators, about 472 species occupy the main honey plants, which include aster, legumes, rosales, labiatae, crucifers (Razanov et al., 2019).

Among the main species of nectar pollinators in Polissya of Ukraine it is necessary to distinguish plants of forest parks, agricultural lands and natural meadows. In particular, the main representatives of nectar-pollinating plants of forest parks of Polissya include linden, maple, acacia; agricultural lands - sunflower, rapeseed and winter rape, buckwheat; natural meadows - dandelion, thyme, white clover, blueweed, clover and others (Ibatullin et al., 2020).

Analysis of the state of nectar-pollinating lands in Polissya of Ukraine shows that among the plants of field crop rotation a promising and powerful source of nectar and pollen is sunflower, which is grown in Ukraine as the main oil crop. In addition, it also provides the beekeeping industry with nectar and pollen, ie raw materials for the production of carbohydrate and protein feed of bees and their products (Nedashkivskiy et al., 2019).

Sunflower belongs to the family Compositae. The duration of its flowering is up to 25 days, and the flowering period - from the second decade of July to the first decade of August. This crop currently occupies the largest sown area among agricultural nectar-pollinating crops in Ukraine, which makes it possible to provide bees in sufficient quantities with both food and raw materials for the production of marketable products (Mazur et al., 2020; Palamarchuk et al., 2021; Rudska, 2021).

Due to the widespread use of beekeeping products as a highly effective food raw material for treatment and prevention, the demand for it is growing rapidly in our country and around the world. At the same time, the requirements for the quality and safety of this product are increasing.

It is known that the quality of beekeeping products depends on the ecological condition of nectar-pollinating lands, which in Ukraine are generally characterized as favorable for the production of high quality beekeeping products, except for some areas that are under high man-made load. It is known that nectar-pollinating lands suffered the highest man-caused load, in particular, some territories of Polissya as a result of the 1986 accident at the Chernobyl nuclear power plant, which caused a certain accumulation of ¹³⁷Cs and ⁹⁰Sr in beekeeping products. At the same time, there were cases of exceeding the permissible levels (DR-1991) for the content of ¹³⁷Cs in honey and pollen in areas where the content of this isotope in the soil was higher than 5 Ci/km². Sometimes there were cases of excess (DR-1991) in beekeeping and ¹³⁷Cs content in soils above 1 Ci/km², especially in bee pollen (pollen) (Dutov et al., 2015; Razanov et al., 2021).

According to the Law of Ukraine "On the Legal Regime of the Territory Contaminated by Radioactive Contamination as a Result of the Chornobyl Accident", the territories, in particular, and agricultural lands where nectar pollinators grow are divided into certain zones:

1. The zone of unconditional resettlement of inhabitants occupies areas with a density of ¹³⁷Cs soil contamination of 15.0 Ci/km² and above, and ⁹⁰Sr - above 3.0 Ci/km².
2. The zone of guaranteed voluntary resettlement of the population includes territories with the level of soil pollution ¹³⁷Cs from 5.0 to 15.0 Ci/km², and ⁹⁰Sr - from 0.15 Ci/km² to 3.0 Ci/km².
3. The zone of enhanced radiological control includes areas with ¹³⁷Cs content in the soil from 1.0 to 5.0 Ci/km², and ⁹⁰Sr - from 0.005 Ci/km² to 0.01 Ci/km² (Law of Ukraine, 1991).

It has been proven that radionuclides that are in soluble soils can migrate into vegetation, reducing their quality and safety. The rate at which ¹³⁷Cs and ⁹⁰Sr move into vegetation depends on a number of factors, including soil type, mineral composition, soil pH, organic matter content, and botanical origin (Frissel et al., 2002; Ji-gen Lu et al., 2006).

Different intensities of radionuclide accumulation both by nectar pollinators and by beekeeping products made from nectar and pollen of these plants were also established. In particular, it was found that the intensive accumulation of ¹³⁷Cs is characterized by such nectar-pollinating plants as thyme, creeping clover, heather, raspberry and others. Honey and pollen produced by bees from the nectar and pollen of these plants were characterized by a higher content of ¹³⁷Cs compared to other nectar pollinators (Razanov et al., 2020).

Peat and peat-swamp soils, which are part of the Polissya ecosystem of Ukraine, are characterized by low sorption properties, on which the transition of ¹³⁷Cs and ⁹⁰Sr to vegetation depends. It was also found that on wet soils the migration of ¹³⁷Cs and ⁹⁰Sr into vegetation is higher compared to soils of moderate moisture (Razanov et al., 2020).

In addition, it should be noted that the activity of movement from soils to plants ¹³⁷Cs and ⁹⁰Sr is their pH. Thus, on soils with high pH of the environment (4.5 - 5.0 units) there is a higher transition of ¹³⁷Cs and ⁹⁰Sr in plants compared to pH 6.0 - 7.0 units (Pryster, 1997).

The danger of ^{137}Cs for the human body, as well as the high demand for bee products, its widespread use for therapeutic and prophylactic purposes among the population, there is a need to control and seek measures to reduce this toxicant in bee products.

One of the measures to increase the safety of beekeeping products, in particular, to reduce ^{137}Cs in it is to prevent the movement of this element in the soil system - nectar pollinators. Chemical reclamation of agricultural lands, which involves liming of soils with high acidity, has a certain effect on the reduction of ^{137}Cs in plants (Camps et al., 2004).

The application of calcium into the soil in the form of lime or defecation mud (waste from sugar factories) leads to the neutralization of the acidity of the soil solution, the displacement of hydrogen ions from the soil complex and its saturation with calcium (Goulding, 2016).

It was found that the application of lime into the soil in an equivalent dose depending on their hydrolytic acidity reduces the transition of ^{137}Cs in crop products up to 2.5 times (Pryster, 1997). The study of ^{137}Cs transition to vegetation in the territory of Polissya of Ukraine was carried out mainly on agricultural crops zoned in these territories (oats, barley, rye, wheat, vetch, peas, corn).

Thirty years after the Chernobyl accident due to climate change, in particular due to rising ambient temperatures, the structural composition of crops has changed somewhat (Bondar, 2020). Thus, recently in the territory of Polissya of Ukraine there is a tendency to increase the area of agricultural land for growing sunflower, which was mainly cultivated until 2000, mainly in the southern steppes of Ukraine. Today, sunflower in the Polissya area is one of the main nectar pollinators among crops (Dikhtiar, 2018).

Due to the high toxicity of ^{137}Cs and the reduction of permissible levels, there is a need to monitor the receipt of this isotope in nectar pollinators and bee products.

Based on this, we monitored the accumulation of ^{137}Cs in beekeeping products and studied the impact of chemical reclamation (application of calcium into the soil as a defecation mud) on its transfer to agricultural nectar-pollinating plants (sunflower) to further predict the entry of this radionuclide into the production of radionuclides, pollen).

MATERIALS AND METHODS

The effectiveness of reducing the migration of ^{137}Cs into agricultural nectar pollinators (sunflower) was carried out in the Narodytsky district of Zhytomyr region, Ukraine.

The research program provided for the study of the specific activity of ^{137}Cs in the soils of agricultural nectar-pollinating lands and in vegetation (vegetative mass) at different stages of their development (stem growth, budding, flowering) for the application of defecation mud into soils. Reclamation of soils included the application of defecation mud in them.

Defecation mud -is the I-class sugar production waste with a content of 60% (Fares et al., 2016). Sunflower cultivation, honey production and bee pollination were carried out in the conditions of four agricultural lands, the distance from each of which was from 8 to 12 km.

The studies were performed on sod-podzolic soils with a contamination density of ^{137}Cs up to 5 Ci/km² and an acidity of 1.6 mg to 2.4 mg-eq/100 g of soil.

In the conditions of agricultural lands № 1 (control) sunflower was grown on soils with ^{137}Cs content - 3640 Bq/kg with hydrolytic acidity of 2.4 mg-eq/100 g of soil without defecation mud.

On agricultural lands №2 (experimental) sunflower was grown on soils with ^{137}Cs content - 3665 Bq/kg and hydrolytic acidity of 2.4 mg-eq/100 g of soil with application of defecation mud at the rate of 6 t/ha.

In the conditions of agricultural lands №3 (control) sunflower cultivation was carried out on soils with ^{137}Cs content of 2824 Bq/kg with hydrolytic acidity of 1.6 mg-eq/100 g of soil without defecation mud.

On agricultural lands №4 (experimental) sunflower cultivation was carried out on soils with ^{137}Cs content 2837 Bq/kg and hydrolytic acidity 1.6 mg-eq/100 g of soil with the application of defecation mud at the rate of 4 t/ha.

The specific activity of ^{137}Cs in soils and vegetative mass of agricultural nectar pollinators was determined using the method of expressive radiometric analysis by gamma radiation of volume and specific activity of Cs radionuclides in water, soil and livestock and crop products.

Soil selection for radiological analysis was performed by envelope method from each field where sunflower was grown.

Selection of vegetative mass of sunflower was carried out by mowing it on an area of 1 m² on 8 sites of each field in different phases of its growth: stem growth (20 days), budding (5 days), flowering (10 days).

The botanical origin of sunflower honey and bee pollen was determined by pollen analysis and the shape of their grains.

The accumulation coefficient of ¹³⁷Cs in different phases of sunflower development and in beekeeping products (honey, bee pollen) was determined by the formula:

$$C_{\text{Accumulation}} = \frac{\text{137Cs content in raw materials}}{\text{137Cs content in soil}}$$

RESULTS AND DISCUSSION

Analysis of the results of studies to study the intensity of ¹³⁷Cs accumulation in the vegetative mass of sunflower (Table 1) showed that the application of defecation mud in the soil helps to reduce the accumulation of this radionuclide in the plant.

Thus, in the phase of sunflower vegetation (stem growth) the specific activity of ¹³⁷Cs during the application of defecation mud in soils with hydrolytic acidity of 2.4 mg decreased by 2.19 times, and the accumulation coefficient by 2.26 times. In the budding phase, the specific activity of ¹³⁷Cs in the vegetative mass of sunflower decreased by 1.99 times, and the accumulation coefficient by 2.04 times. Whereas in the sunflower flowering phase the specific activity of ¹³⁷Cs and the coefficient of accumulation in the vegetative mass decreased by 2.3 times and 2.4 times, respectively.

Table 1. The intensity of the accumulation of ¹³⁷Cs in the vegetative mass of sunflower for the application of defecation mud into the soil

Development phase	Indexes	Agricultural land			
		№1 (without application of defecation mud)	№2 (with application of defecation mud)	№3 (without application of defecation mud)	№4 (with application of defecation mud)
Stem growth	Specific activity of ¹³⁷ Cs in soil, Bq/kg	3640±3.2	3665±7.8	2824±1.9	2837±2.7
	The specific activity of ¹³⁷ Cs in the plant, Bq/kg	191.4±4.7	87.1±1.2	152.1±1.8	81.7±0.7
	Accumulation coefficient	0.052	0.023	0.053	0.028
Budding	Specific activity of ¹³⁷ Cs in soil, Bq/kg	3640±1.2	3665±7.8	2824±4.1	2837±2.7
	The specific activity of ¹³⁷ Cs in the plant, Bq/kg	157.2±1.4	78.8±0.8	143.5±2.1	73.7±1.3
	Accumulation coefficient	0.043	0.021	0.050	0.033
Flowering	Specific activity of ¹³⁷ Cs in soil, Bq/kg	3640±3.2	3665±7.8	2824±4.1	2837±2.7
	The specific activity of ¹³⁷ Cs in the plant Bq/kg	150.3±2.7	65.1±0.6	128±3.4	59.8±2.7
	Accumulation coefficient	0.041	0.017	0.045	0.021

A similar trend was observed for the application of defecation mud in soils with a hydrolytic acidity of 1.6 mg. In particular, when defecation mud were applied into soils with hydrolytic acidity of 1.6 mg in the vegetative mass of sunflower in the stem growth phase, the specific activity of ^{137}Cs was 1.86 times lower and the accumulation coefficient was 1.89 times. Whereas in the budding and flowering phase of sunflower, the specific activity of ^{137}Cs during the application of defecation mud into the soil was 1.94 times and 1.92 times lower, and the accumulation coefficient was 2.14 and 2.14 times, respectively, compared to the variant without fertilizer.




Figure 1. Comparative assessment of ^{137}Cs reduction in the vegetative mass of sunflower for the application of defecation mud in soils of different acidity

Characterizing the effectiveness of reducing ^{137}Cs in the vegetative mass of sunflower (Fig. 1), it should be noted that the highest result was achieved by making defecation mud in soils with hydrolytic acidity of 2.4 mg compared to soils with hydrolytic acidity of 1.6 mg.

In particular, the rate of decrease in ^{137}Cs in the vegetative mass of sunflower in the growth phase (stem growth, budding, flowering) for the application of defecation mud in soils with hydrolytic acidity of 2.4 mg was higher by 0.33 percentage points (pp), 0.05 pp and 0.16 pp respectively compared to the hydrolytic acidity of 1.6 mg.

Table 2. The intensity of the accumulation ^{137}Cs in beekeeping products made from nectar and pollen of sunflower for the application of defecation mud into the soil

Agricultural lands and research options	Beekeeping products					
	Honey			Bee pollen		
	Specific activity in soil ^{137}Cs , Bq/kg	The specific activity of ^{137}Cs in honey Bq/kg	$C_{\text{accumulation}}$	Specific activity in soil ^{137}Cs , Bq/kg	The specific activity of ^{137}Cs in bee pollen Bq/kg	$C_{\text{accumulation}}$
№1 – without application of defecation mud (control)	3640±5.1	32.3±0.7	0.008	3640±5.1	81.5±1.2	0.022
№2 – with application of defecation mud (experimental)	3665±2.1	17.2±0.34	0.004	3665±2.1	27.2±0.74	0.007
№3 – without application of defecation mud (control)	2824±1.4	30.4±0.95	0.01	2824±1.4	70.3±2.1	0.024
№4 – with application of defecation mud (experimental)	2837±1.5	18.1±1.5	0.006	2837±1.5	28.1±1.7	0.009

We have found out changes in the intake of ¹³⁷Cs in beekeeping products during liming of acid soils of agricultural lands for growing sunflower, the nectar and pollen of which are raw materials for the production of honey by bees and bee pollen (Table 2).

Thus, the specific activity and accumulation coefficient of ¹³⁷Cs in honey produced by bees from sunflower nectar during its cultivation when making defecation mud in soils with hydrolytic acidity of 2.4 mg, was lower by 1.87 times and 2.0 times compared to the option without making defecation mud.

When defecation mud were introduced into soils with hydrolytic acidity of 1.6 mg, the specific activity of ¹³⁷Cs in honey produced by bees from sunflower nectar was 1.68 times lower and the accumulation coefficient was 1.66 times lower than in the non-defecation mudive version.

A similar tendency to decrease the specific activity and accumulation coefficient of ¹³⁷Cs was observed in bee pollen produced by bees from sunflower pollen for defecation in acidic soils of agricultural lands.

In particular, when the defecation mud was introduced into soils with hydrolytic acidity of 2.4 mg, the specific activity of ¹³⁷Cs in bee pollen made by bees from sunflower pollen was 3.0 times lower and the accumulation coefficient was 3.1 times lower than in the non-defecation mudive version. .

When defecation mud were introduced into soils with a hydrolytic acidity of 1.6 mg, the specific activity of ¹³⁷Cs in bee pollination was 2.5 times lower, and the accumulation coefficient was 2.6 times lower than in the variant without defecation.

In addition, it should be noted that the specific activity of ¹³⁷Cs in honey and bee pollen for the application of defecation mud in soils with hydrolytic acidity of 2.4 mg was lower than DR-2000 g 11.6 times and 7.3 times, respectively, whereas in the version without these defecation mud were 6.2 and 2.4 times higher in the soil.

The specific activity of ¹³⁷Cs in honey and bee pollen produced by bees from the nectar of sunflower pollen when growing it for defecation in soils with hydrolytic acidity of 1.6 mg was 11.0 times lower than DR-2000 and 7.1 times, respectively, while in the version without introducing defecation mud into the soil (control), these figures were between 6.5 and 2.8 times.




Figure 2. Comparative assessment of the reduction of ¹³⁷Cs movement in beekeeping products for the application of defecation mud in soils of different acidity

There was also a higher efficiency of ¹³⁷Cs reduction in beekeeping products with the application of defecation mud in soils with hydrolytic acidity of 2.4 mg-eq/100 g compared to hydrolytic acidity of 1.6 mg-eq/100 g (Fig. 2).

Thus, the application of defecation mud into soils with hydrolytic acidity of 2.4 mg showed a decrease in ¹³⁷Cs in honey by 0.19 percentage points, and in bee pollination by 0.5 percentage points compared with hydrolytic acidity of 1.6 mg.

Thus, the results of research revealed that reclamation of agricultural nectar pollinators (application into the soil of defecation mud) reduces the accumulation of ¹³⁷Cs in the vegetative mass of sunflower, honey and bee pollen produced by bees from nectar and pollen.

CONCLUSION

- The specific activity of ^{137}Cs in honey and bee pollen produced by bees from nectar and pollen of sunflower grown on agricultural land with the content of this radionuclide in soils from 2824 Bq/kg to 2665 Bq/kg, did not exceed 200 DR, 2006 Bq/kg.
- Reclamation of radioactively contaminated soils with hydrolytic acidity from 1.6 mg to 2.4 mg-eq/100 g of soil, in particular, the application of defecation mud in them at a rate of 4 t/ha to 6 t/ha reduced the specific activity and accumulation rate ^{137}Cs in the vegetative mass of sunflower, honey and bee pollen, made by bees from the nectar and pollen of this plant.
- Higher efficiency of reduction of specific activity and accumulation coefficient of ^{137}Cs in honey and bee pollen produced from nectar and pollen of sunflower was characterized by defecation mud application in soils with hydrolytic acidity of 2.4 mg-eq/100 g of soil compared to 1.6 mg-eq/100 g of soil.

REFERENCES

1. Bondar, I.M. 2020. The use of agricultural machinery in the technology of sunflower cultivation in the Polissya region. Bulletin of the Petro Vasylenko Kharkiv National Technical University of Agriculture. 209: 15-25;
2. Camps, M., Rigol, A., Hillier, S., Vidal, M., Rauret, G. 2004. Quantitative assessment of the effects of agricultural practices designed to reduce ^{137}Cs and ^{90}Sr soil-plant transfer in meadows. Science of the total environment. 332 (1-3): 23-38;
3. Dikhtiar, O.O. 2018. The quality of the sunflower honey obtained under the conditions of radiologically contaminated agronomic landscape Polissia region. Scientific Bulletin of the National University of Life and Environmental Sciences of Ukraine. 289: 163-170;
4. Dutov, O.I., Landin, V.P., Melnychuk, A.O., Hrynyk, O.I. 2015. Radiation environmental aspects of contaminated land in the remote period after the accident at the Chernobyl nuclear power plant. Agroecological journal. 1: 115-120;
5. Fares, K., Saadaoui, N. and RıZina, Q. 2016. Sugar beet lime sludge composts as organic fertilizers. Acta Horticulturae. 1146: 165-170;
6. Frissel, M.J., Deb, D.L., Fathony, M., Lin, Y.M., Mollah, A.S., Ngo, N.T., Othman, I., Robison, W.L., SkarlouAlexiou, V., Topcuoglu, S., Twining, J.R., Uchida, S., Wasserman, M.A. 2002. Generic values for soil-to-plant transfer factor of radiocesium. Journal of Environmental Radioactivity. 58: 113-128;
7. Goulding, K. W. T. 2016. Soil acidification and the importance of liming agricultural soils with particular reference to the United Kingdom. Soil Use and Management. 32: 390-399;
8. Ibatullin, I.I., Razanov, S.F., Nedashkivskyi, V.M. 2020. Influence of soy flour and its processed products on the essential amino acids content in the bees body. Veterinary and Agricultural Sciences. 3 (3): 51-55;
9. Ji-gen Lu, Yanjun Huang, Fang Li, Lihua Wang, Shuzhen Li, Yuanfu Hsia. 2006. The investigation of ^{137}Cs and ^{90}Sr background radiation levels in soil and plant around Tianwan NPP, China. Journal of Environmental Radioactivity. 90: 89-99;
10. Law of Ukraine. 1991. «On the legal regime of the territory exposed to radioactive contamination as a result of the Chernobyl disaster»;
11. Mazur, V.A., Didur, I.M., Tsyhanskyi, V.I., Malamura, S.V. 2020. Formation of productivity of sunflower hybrids depending on the level of fertilizer and moisture conditions. Agriculture and forestry. 19: 208-220;
12. Nedashkivskyi, V.M., Kovka, N.O. 2019. Duration and periods of flowering of the main nectar pollinators in the conditions of the Forest-Steppe of the Right Bank. Livestock of Ukraine. 4: 36-39;
13. Nicolson, S.W. 2011. Bee Food: The Chemistry and Nutritional Value of Nectar, Pollen and Mixtures of the Two. African Zoology. 46 (2): 197-204;
14. Palamarchuk, V.D., Pidlubnyi, V.F. 2021. Productivity of sunflower hybrids depending on the elements of the growing technology. Agriculture and forestry. 3 (22): 29-44;

15. Pryster, B.S. 1997. Handbook for radiological services of the Ministry of Agriculture of Ukraine. Kyiv. 175 p;
16. Razanov, S.F., Nedashkivskyi, V.M. 2019. Nectar pollinator conveyor of bees in the Forest-Steppe of the Right Bank. Livestock of Ukraine. 2: 11-14;
17. Razanov, S.F., Ogorodnichuk, G.M., Komynar, M.F. 2021. Effect of soil treatment on ¹³⁷Cs accumulation in flower pollen and in its processing products by honey bees. Agriculture and forestry. 3 (22): 161-173;
18. Razanov, S.F., Shevchuk, V.V., Komynar, M.F. 2020. Accumulation of ¹³⁷Cs in honey produced by bees from nectar of agricultural honeybear in the northern Polisia. Agriculture and forestry. 19: 148-158;
19. Rudska, N.O. 2021. Influence of technological techniques and improvement of the system of protection of sunflower crops from weeds. Colloquium-journal. 16 (103): 22-30;