



Founder, Editorial and Publisher
POLISSIA NATIONAL UNIVERSITY

e-ISSN 2709-8877

ISSN: 2663-2144

SCIENTIFIC

H O R I Z O N S



НАУКОВІ ГОРИЗОНТИ

Засновник, редакція, видавець:
Поліський національний університет

Рік заснування: 1998

*Рекомендовано до друку та поширення
через мережу Інтернет Вченою радою
Поліського національного університету
(протокол № х від хх січня 2023 року)*

**Свідоцтво про державну реєстрацію
друкованого засобу масової інформації
серії KB № 24997-14937 ПР**

Науковий журнал включено до категорії А Переліку наукових фахових видань України, у яких можуть публікуватися результати дисертаційних робіт на здобуття наукових ступенів доктора та кандидата ветеринарних, економічних, сільськогосподарських і технічних наук зі спеціальностей: 051 – Економіка; 071 – Облік і оподаткування; 072 – Фінанси, банківська справа та страхування; 073 – Менеджмент; 075 – Маркетинг; 076 – Підприємництво, торгівля та біржова діяльність; 101 – Екологія; 133 – Галузеве машинобудування; 201 – Агрономія; 202 – Захист і карантин рослин; 203 – Садівництво та виноградарство; 204 – Технологія виробництва і переробки продукції тваринництва; 205 – Лісове господарство; 206 – Садово-паркове господарство; 208 – Агроінженерія; 211 – Ветеринарна медицина (наказ МОН України № 1166 від 23 грудня 2022 р.)

Журнал представлено у міжнародних наукометричних базах даних, репозитаріях та пошукових системах: Інституційний репозитарій Поліського національного університету, AGRICOLA, CAB Abstracts and Global Health (CABI), Open Academic Journals Index (OAJI), Scopus, Index Copernicus, Реєстр наукових видань України, Національна бібліотека України імені В. І. Вернадського, Crossref, Directory of Open Access Journals (DOAJ)

Адреса редакції:
Поліський національний університет
б-р Старий, 7, м. Житомир, 10008, Україна
E-mail: info@sciencehorizon.com.ua
<https://sciencehorizon.com.ua/uk>

SCIENTIFIC HORIZONS

Founder, Editorial and Publisher:

Polissia National University

Year of foundation: 1998

*Recommended for printing and distribution
via the Internet by the Academic Council
of Polissia National University
(Minutes No. x of January xx, 2023)*

Certificate of state registration

print mass media

Series KV No. 24997-14937 PR

The scientific journal is included in category A of the List of scientific professional publications of Ukraine, in which the results of dissertations for the degree of doctor and candidate of veterinary, economic, agricultural, and technical sciences can be published in the following specialties: 051 – Economy; 071 – Accounting and Taxation; 072 – Finance, Banking and Insurance; 073 – Management; 075 – Marketing; 076 – Entrepreneurship, Trade and Exchange Activities; 101 – Ecology; 133 – Sectoral Engineering; 201 – Agronomy; 202 – Plant Protection and Quarantine; 203 – Horticulture and Viticulture; 204 – Technology of Production and Processing of Livestock Products; 205 – Forestry; 206 – Park and Gardening Management; 208 – Agricultural Engineering; 211 – Veterinary Medicine (Order of the Ministry of Education and Science of Ukraine No. 1166 of December 23, 2022)

The journal is presented international scientometric databases, repositories

and scientific systems: Institutional Repository of Polissia National University, AGRICOLA, CAB Abstracts and Global Health (CABI), Open Academic Journals Index (OAJI), Scopus, Index Copernicus, Register of Scientific Publications of Ukraine, Vernadsky National Library of Ukraine, Crossref, Directory of Open Access Journals (DOAJ)

Editors office address:

Polissia National University
10008, 7 Staryi Blvd., Zhytomyr, Ukraine
E-mail: info@sciencehorizon.com.ua
<https://sciencehorizon.com.ua/en>

НАУКОВІ ГОРИЗОНТИ

Том 26, № 2 2023

НАУКОВИЙ ЖУРНАЛ
Засновано 12 березня 1998 р.

Періодичність випуску: дванадцять разів на рік

Редакційна колегія

Головний редактор Олег Васильович Скидан	ректор, д-р екон. наук, професор, Поліський національний університет, Україна
Заступник головного редактора Людмила Романчук	д-р с.-г. наук, професор, Поліський національний університет, Україна
Національні члени редколегії	
Савелій Кухарець	д-р техн. наук, професор, Поліський національний університет, Україна
Іван Грабар	д-р техн. наук, професор, Поліський національний університет, Україна
Ярослав Ярош	д-р техн. наук, професор, Поліський національний університет, Україна
Тетяна Федонюк	д-р с.-г. наук, професор, Поліський національний університет, Україна
Наталія Сорока	д-р вет. наук, професор, Національний університет біоресурсів і природокористування України, Україна
Руслана Ставецька	д-р с.-г. наук, професор, Білоцерківський національний аграрний університет, Україна
Анастасія Зимароєва	канд. біол. наук, доцент, Поліський національний університет, Україна
Людмила Чижевська	д-р екон. наук, професор, Державний університет «Житомирська політехніка», Україна
Віталій Данкевич	д-р екон. наук, професор, Поліський національний університет, Україна
Ельчин Алієв	д-р техн. наук., професор, Дніпровський державний аграрно-економічний університет, Україна
Олександр Галатюк	д-р вет. наук, професор, Поліський національний університет, Україна
Валентина Гамаюнова	д-р с.-г. наук, професор, Миколаївський національний аграрний університет, Україна
Леонід Горальський	д-р вет. наук, професор, Житомирський державний університет імені Івана Франка, Поліський національний університет, Україна
Дмитро Дерев'янюк	д-р техн. наук, професор, Поліський національний університет, Україна
Олександр Жуков	д-р біол. наук, професор, Мелітопольський державний педагогічний університет імені Богдана Хмельницького, Україна
Ірина Іванова	канд. с.-г. наук, доцент Таврійський державний агротехнологічний університет ім. Д. Моторного, Україна
Олександр Кочук-Ященко	канд. с.-г. наук, доцент, Поліський національний університет, Україна
Дмитро Кучер	канд. с.-г. наук, доцент, Поліський національний університет, Україна
Федір Марков	канд. с.-г. наук, доцент, Поліський національний університет, Україна
Олена Марковська	д-р с.-г. наук, старший науковий співробітник, Херсонський державний аграрно-економічний університет, Україна
Андрій Михайлов	д-р екон. наук, професор, Сумський національний аграрний університет, Україна
Валентин Москалець	д-р с.-г. наук, доцент, Інститут садівництва НААН, Україна
Віктор Пазич	канд. с.-г. наук, доцент, Поліський національний університет, Україна
Людмила Тарасович	канд. екон. наук, доцент, Поліський національний університет, Україна
Тетяна Тимошук	канд. с.-г. наук, доцент, Поліський національний університет, Україна
Наталія Цивенкова	канд. техн. наук, доцент, Національний університет біоресурсів і природокористування України, Поліський національний університет, Україна
Марина Швець	канд. біол. наук, старший викладач Поліський національний університет, Україна

Тетяна Зінчук	д-р екон. наук, професор, Поліський національний університет, Україна
Геннадій Голуб	д-р техн. наук, професор, Національний університет біоресурсів і природокористування України, Україна
Міжнародні члени редколегії	
Юліус Раманаускас	д-р наук, професор, Клайпедський університет, Литва
Інна Левкович	д-р наук, старший науковий співробітник, Лейбніцький інститут розвитку сільського господарства у країнах з перехідною економікою, Німеччина
Егідіюс Сараускіс	д-р техн. наук, професор, Інститут сільськогосподарської техніки та безпеки університету Вітовта Великого, Литва
Ришард Пукала	канд. екон. наук, Державний Техніко-Економічний Університет ім. кс. Броніслава Маркевича в Ярославі, Польща
Самат Танібергенов	вчений секретар, Казахський науково-дослідний інститут ґрунтознавства і агрохімії ім. У.У. Успанова, Республіка Казахстан
Саліх Ташпулатов	д-р техн. наук, професор, Ташкентський інститут текстильної та легкої промисловості, Республіка Узбекистан

SCIENTIFIC HORIZONS

Vol. 26, No. 2
2023

SCIENTIFIC JOURNAL
Year of establishment: Since March 1998.
Publication frequency: Twelve times a year

Editorial Board

Editor-in-Chief

Oleh V. Skydan Rector, Full Doctor in Economic Sciences, Professor, Polissia National University, Ukraine

Deputy Editor-in-Chief

Ludmila Romantschuk Full Doctor in Agricultural Sciences, Professor, Polissia National University, Ukraine

National Members of the Editorial Board

Savelli Kukharets Full Doctor in Engineering Sciences, Professor, Polissia National University, Ukraine

Ivan Grabar Full Doctor in Engineering Sciences, Professor, Polissia National University, Ukraine

Yaroslav Yarosh Full Doctor in Engineering Sciences, Professor, Polissia National University, Ukraine

Tatiana Fedoniuk Full Doctor in Agricultural Sciences, Professor, Polissia National University, Ukraine

Natalia Soroka Full Doctor in Veterinary Sciences, Professor, National University of Life and Environmental Sciences of Ukraine, Ukraine

Ruslana Stavetska Full Doctor in Agricultural Sciences, Professor, Bila Tserkva National Agrarian University, Ukraine

Anastasiia Zymarioieva PhD in Biological Sciences, Associate Professor, Polissia National University, Ukraine

Lyudmyla Chyzhevska Full Doctor in Economic Sciences, Professor, Zhytomyr Polytechnic State University, Ukraine

Vitalii Dankevych Full Doctor in Economic Sciences, Professor, Polissia National University, Ukraine

Elchin Aliiev Full Doctor in Engineering Sciences, Professor, Dnipro State Agrarian and Economic University, Ukraine

Olexandr Galatiuk Full Doctor in Veterinary Sciences, Professor, Polissia National University, Ukraine

Valentina Hamaiunova Full Doctor in Agricultural Sciences, Professor, Mykolayiv National Agrarian University, Ukraine

Leonid Goralskiy Full Doctor in Veterinary Sciences, Professor, Zhytomyr Ivan Franko State University, Polissia National University, Ukraine

Dmytro Derevjanko Full Doctor in Engineering Sciences, Professor, Polissia National University, Ukraine

Olexander Zhukov Full Doctor in Biological Sciences, Professor, Bogdan Khmelnytsky Melitopol State Pedagogical University, Ukraine

Iryna Ivanova PhD in Agricultural Sciences, Associate Professor, Dmytro Motornyi Tavria State Agrotechnological University, Ukraine

Oleksandr Kochuk-Yashchenko PhD in Agricultural Sciences, Associate Professor, Polissia National University, Ukraine

Dmytro Kucher PhD in Agricultural Sciences, Associate Professor, Polissia National University, Ukraine

Fedir Markov PhD in Agricultural Sciences, Associate Professor, Polissia National University, Ukraine

Olena Markovska Full Doctor in Agricultural Sciences, Senior Research Scientist, Kherson State Agrarian and Economic University, Ukraine

Andrii Mykhailov Full Doctor in Economic Sciences, Professor, Sumy National Agrarian University, Ukraine

Valentyn Moskalets Full Doctor in Agricultural Sciences, Associate Professor, Institute of Horticulture of the National Academy of Agrarian Sciences, Ukraine

Viktor Pazych PhD in Agricultural Sciences, Associate Professor, Polissia National University, Ukraine

Ludmyla Tarasovych PhD in Economic Sciences, Associate Professor, Polissia National University, Ukraine

Tetiana Tymoshchuk
Natalia Tsyvenkova

PhD in Agricultural Sciences, Associate Professor, Polissia National University, Ukraine
PhD in Engineering Sciences, Associate Professor, National University of Life and Environmental Sciences of Ukraine, Polissia National University, Ukraine

Marina Shvets
Tetyana Zinchuk
Gennadii Golub

PhD in Biological Sciences, Senior Teacher, Polissia National University, Ukraine
Full Doctor in Economic Sciences, Professor, Polissia National University, Ukraine
Full Doctor in Engineering Sciences, Professor, National University of Life and Environmental Sciences of Ukraine, Ukraine

**International members
of the Editorial Board**

Julius Ramanauskas
Inna Levkovich

Dr. Habil., Professor, Klaipeda University, Lithuania
Dr. Habil., Senior Research Scientist, Leibniz Institute of Agricultural Development in Transition Economies (IAMO), Germany

Egidijus Sarauskis

Full Doctor in Engineering Sciences, Professor, Institute of Agricultural Engineering and Safety of Vytautas Magnus university (VMU), Lithuania

Ryszard Pukala

PhD in Economic Sciences, The Bronisław Markiewicz State University of Technology and Economics in Jarosław, Poland

Samat Tanirbergenov

Scientific Secretary, U.U. Uspanov Kazakh Research Institute of Soil Science and Agrochemistry, Republic of Kazakhstan

Salikh Tashpulatov

Full Doctor in Engineering Sciences, Professor, Tashkent Institute of Textile and Light Industry, Republic of Uzbekistan

ЗМІСТ

Л. О. Шевчик, І. М. Грод, Н. Я. Кравець, Я. С. Стравський, М. А. Крижановська Етологічна адаптивність домашніх тварин на прикладі різних порід kota свійського (<i>Felis silvestris catus</i> L., 1758).....	9
Г. В. Миронова, Т. М. Тимошук, О. В. Волошина, О. В. Мазур, О. В. Мазур Формування урожайності насінневої картоплі залежно від елементів технології вирощування.....	19
С. Атанасов, Б. Харізанова-Петрова, Р. Петрова Колір листя томатів як показник вологості ґрунту з використанням методів машинного навчання.....	31
Ю. Е. Клечковський, Н. Т. Могилюк, К. А. Шматковська, І. М. Сидорова, С. М. Кубрак Стемфіліоз цибулі ріпчастої на півдні України	43
О. В. Мельник, Н. Г. Духіна, В. І. Пастухов, О. Л. Семенченко, Є. М. Ільїнова Фізіолого-біохімічне обґрунтування двоврожайної культури картоплі (<i>Solanum tuberosum</i> L.).....	54
М. Б. Грабовський, О. В. Мостипан, Ю. В. Федорук, Л. А. Козак, М. В. Остренко Формування урожайності зерна та якісних показників сої під впливом фунгіцидного захисту.....	66
В. Г. Кургак, Л. П. Коломієць, О. М. Корягін, Л. М. Красюк, Р. М. Кулик Мінеральний склад біомаси люцернового, люцерно-злакових і злакового кормових агрофітоценозів.....	77
Н. М. Вознюк, В. П. Скиба, О. А. Ліхо, З. З. Собко, Т. В. Клименко Прогнозування адаптивності теплолюбних культур до змін клімату в умовах України.....	87
Х. Кічай Екологічні фактори, що впливають на поширення класу Chilopoda групи Mugiapoda	103
М. Ісмаїл-заде Аналіз ефективності впровадження фізико-економічної теорії в економічну діяльність Азербайджану.....	112
М. В. Дубініна, Н. В. Потриваєва, О. М. Вишневська, О. М. Усикова, В. В. Кузьома Інституційні аспекти демонстрації обліку трансакційних витрат в сільськогосподарських організаціях.....	124
ОГЛЯДОВА СТАТТЯ	
О. М. Семенченко, Ю. Б. Добровольський, П. О. Толок, А. П. Онофрійчук, В. П. Онофрійчук Енергетична та екологічна безпека Європейського Союзу в умовах нестабільної військової та економічної політики росії.....	135

CONTENTS

L. Shevchyk, I. Grod, N. Kravets, Y. Stravskyy, M. Kryzhanovska

Behavioural adaptation of domestic animals on the example of different breeds of domestic cat
(*Felis silvestris catus* L., 1758)9

H. Myronova, T. Tymoshchuk, O. Voloshyna, O. Mazur, O. Mazur

Formation of seed potato yield depending on the elements of cultivation technology19

S. Atanasov, B. Harizanova-Petrova, R. Petrova

Tomato leaf colour as predictor of soil moisture level using machine learning techniques31

Yu. Klechkovskyi, N. Mogilyuk, K. Shmatkovska, I. Sidorova, S. Kubrak

Onion stemphiliosis in the South of Ukraine43

O. Melnyk, N. Dukhina, V. Pastukhov, O. Semenchenko, Ye. Ilinova

Physiological and biochemical justification of second-crop potatoes (*Solanum tuberosum* L.).....54

M. Grabovskyi, O. Mostypan, Y. Fedoruk, L. Kozak, M. Ostrenko

Formation of grain yield and quality indicators of soybeans under the influence of fungicidal protection66

V. Kurhak, L. Kolomiets, O. Koryagin, L. Krasiuk, R. Kulyk

Mineral composition of alfalfa, alfalfa-cereal, and cereal feed agrophytocenoses biomass77

N. Vozniuk, V. Skyba, L. Olena, S. Zoriana, K. Tetiana

Forecasting the adaptability of heat-loving crops to climate change in Ukraine87

H. Kicaj

Ecological Factors Affecting the Spread of Class Chilopoda of the Group Myriapoda103

M. Ismayil-zada

Analysis of physical economic theory implementation efficiency in the economic activity of Azerbaijan.....112

M. Dubinina, N. Potryvaieva, O. Vyshnevskaya, O. Usykova, V. Kuzoma

Institutional aspects of transactional spending accounting demonstration in agricultural organisations124

REVIEW ARTICLE

O. Semenenko, U. Dobrovolskyi, P. Tolok, A. Onofriichuk, V. Onofriichuk

Energy and environmental security of the European Union
in the context of Russia's unstable military and economic policy.....135



UDC 631.559:633.49:631.5

DOI: 10.48077/scihor.26(2).2023.19-30

Formation of seed potato yield depending on the elements of cultivation technology

Hanna Myronova

Postgraduate Student

Vinnitsia National Agrarian University
21008, 3 Sonyachna Str., Vinnitsia, Ukraine
<https://orcid.org/0000-0001-8401-6138>

Tetiana Tymoshchuk*

Candidate of Agricultural Sciences, Associate Professor
Polissia National University
10008, 7 Staryi Blvd., Zhytomyr, Ukraine
<https://orcid.org/0000-0001-8980-7334>

Oksana Voloshyna

Candidate of Philological Sciences, Associate Professor
Vinnitsia National Agrarian University
21008, 3 Sonyachna Str., Vinnitsia, Ukraine
<https://orcid.org/0000-0002-7679-9555>

Olena Mazur

Candidate of Agricultural Sciences, Associate Professor
Vinnitsia National Agrarian University
21008, 3 Sonyachna Str., Vinnitsia, Ukraine
<https://orcid.org/0000-0003-0132-7470>

Oleksandr Mazur

Candidate of Agricultural Sciences, Associate Professor
Vinnitsia National Agrarian University
21008, 3 Sonyachna Str., Vinnitsia, Ukraine
<https://orcid.org/0000-0002-2237-5116>

Article's History:

Received: 26.12.2022

Revised: 18.02.2023

Accepted: 10.03.2023

Suggested Citation:

Myronova, H., Tymoshchuk, T., Voloshyna, O., Mazur, O., & Mazur, O. (2023). Formation of seed potato yield depending on the elements of cultivation technology. *Scientific Horizons*, 26(2), 19-30.

Abstract. It is possible to realise the maximum genetic potential of modern potato varieties by improving the elements of agricultural technology, in particular, optimising plant nutrition, which determines the relevance of this study. The purpose of the study was to substantiate the specific features of potato productivity formation depending on the elements of growing technology in Forest-Steppe conditions. Field studies were conducted during 2019-2021 on chernozem soils. The features of growth and development of potato varieties of various ripeness groups Laperla, Granada, and Memphis in Forest-Steppe conditions are examined. It is established that the yield of potato seed tubers of the varieties under study varies depending on the dose and method of fertilisation, the fraction of planting tubers, and varietal characteristics. The highest indicators of



individual productivity of potato plants of the Laperla, Granada, and Memphis varieties were formed when locally applied in rows $N_{45}R_{45}K_{45}$ against the background of semi-rotted manure, phosphorus-potassium fertiliser, and the use of seed tubers with a fraction of >60 mm. As a result of the action of the factors under study, the number of stems in the bush and under it increased. A strong direct correlation was established between the number of stems in the bush and the number of tubers ($r=0.78$), with a coefficient of determination of 61.0%. The maximum yield of potato tubers of the examined potato varieties (29.7-41.1 t/ha) was obtained by local application of $N_{45}R_{45}K_{45}$ against the background of semi-rotted manure and phosphorus-potassium fertiliser, depending on the size of seed tubers by the largest transverse diameter. It was identified that the use of tubers with a fraction of >60 mm for planting the potato varieties under study is irrational since the consumption of seed material exceeds the yield growth indicators. It was determined that Laperla, Granada, and Memphis tubers with a fraction of 28-60 mm are optimal for planting potato varieties of different ripeness groups. The research results can be used to adjust the elements of agricultural technology of seed potatoes, obtain high-quality planting material, and conduct profitable agribusiness

Keywords: fertiliser; yield structure; individual plant productivity; variety; tuber size; tuber fraction

INTRODUCTION

Potatoes are one of the most widespread agricultural crops in the world, which ranks fourth after rice, wheat, and corn in terms of value and importance for the world's population and ensuring food security. Potato cultivation in many countries is conducted not only by large agricultural enterprises but also by small farms. This popularity of potato commercialisation is explained by its balance as a human food product, animal feed, and raw materials for the processing industry. Modern potato varieties and hybrids have a high potential for adaptability and productivity, which is the basis for obtaining stable high yields with valuable consumer, technical, and economic properties. Currently, many potato varieties have been imported to Ukraine from other countries, which are not yet fully adapted to the new growing conditions. Therefore, studies of the response of modern varieties to fertiliser optimisation depending on the fraction of seed tubers are relevant and of practical interest.

Balanced use of nitrogen, phosphorus and potash fertilisers is an important factor for maximising crop yields and achieving the desired quality. The study by researchers (Kyryliuk *et al.*, 2020; Biliavska *et al.*, 2021; Kotelnyska *et al.*, 2021) confirmed the improvement of the nutrient regime of the arable soil layer, optimisation of its agrophysical and water-physical indicators by applying organic and mineral fertilisers, which are mineralised by soil microorganisms and provide plants with available nutrients. The positive effect of optimising potato fertiliser on the yield and quality of tubers was noted in the works of many researchers. According to O. Iskakova and V. Gamajunova (2021), application of mineral fertilisers in autumn ($N_{32}P_{32}K_{32}$), before planting ($N_{48}P_{48}K_{48}$) and three-time foliar top dressing (Plantafol, 6 kg/ha) at the beginning of budding in 8-10 days ensures the formation of tuber yield at the level of 37-39 t/ha. The authors also investigated the effect of optimising potato nutrition on reducing the content of dry substances and starch in tubers and increasing

the content of ascorbic acid. According to M. Polishchuk (2021a) application of organic and mineral fertilisers (manure 40 t/ha+ $N_{90}P_{90}K_{90}$ and manure 40 t/ha+ $N_{120}P_{120}K_{120}$) and potato planting rates of 50 thousand tubers/ha ensures the formation of the highest yield – 29.3 and 30.5 t/ha, respectively.

Now there is a discrepancy between the norms of planting potatoes and the yield of tubers. Thus, the consumption rates of planting material range from 3.0 to 5 t/ha, and the yield is only 11.0-16.0 t/ha, depending on the year of cultivation (Polishchuk, 2021b). Thus, the ratio between the yield and the planting rate (reproduction rate) is only 2.5-3.0, which makes the cultivation of this crop economically impractical. The correct choice of the size of planted potato tubers and compliance with scientifically based planting density are the crucial factors in solving this problem (Kumar & Aulakh, 2022).

A study by M.J. Sadawarti *et al.* (2021) established the effect of seed tuber size on the overall yield and output of commercial tubers. In the paper of S. Ebrahim *et al.* (2018), a research on the effect of seed tuber size (fractions 25-54, 35-45, and 46-55 mm) on the productivity of three potato varieties Gudenie, Jalene, and Kellacho in Ethiopia is presented. It is established that planting potatoes with large seed tubers ensures the formation of a maximum yield of 42.9 t/ha.

Studies conducted in the northwestern part of India are devoted to the examination of the influence of seed tuber size (25-55 mm, 35-45 mm, 45-55 mm) on potato tuber contamination and yield (Kumar *et al.*, 2021). Researchers identified that planting potatoes with medium (35-45 mm) and large (45-55 mm) seed tubers reduces the number and biomass of weeds, improves seed productivity and increases potato yield to 42.6 and 42.9 t/ha, respectively. However, they identified that it is more reasonable to plant tubers of medium size (35-45 mm).

The fraction of seed tubers affects the stem-forming ability of potato varieties (Ilchuk & Ilchuk, 2021). The maximum potato stalk was formed by varieties Kimmeriia (330

and 342 thousand stems/ha, respectively) and Shchedryk (268 and 375 thousand stems/ha, respectively) when planted with tubers with a fraction of ≥ 45 mm and ≥ 60 mm.

Consequently, the production of high-quality potato seed tubers is an important issue in traditional potato growing regions and should be improved by optimising agricultural technologies. Considering the above, the solution to the problem of growing potato seed material becomes relevant and requires the optimisation of elements of cultivation technology. The purpose of the study was to establish the dependence of the formation of the number of stems, the number of tubers under the bush, the mass of tubers from one bush on fertiliser, the fraction of seed material, and the varietal characteristics of potatoes.

MATERIALS AND METHODS

Research on the study of technology elements for potato productivity was conducted during 2019-2021 in the conditions of the farm "Olvia-S" (Sopyn village, Vinnytsia district, Vinnytsia region). The territory of the farm is located in the Forest-Steppe zone. The soil cover of the territory where the study was conducted is represented by deep low-humus medium-loamy chernozems. The arable layer (0-30 cm) of the soil of the experimental site has the following agrochemical indicators: humus content (according to Tyurin) 4.42%, easily hydrolysed nitrogen (according to Kornfeld) – 12.0 mg/100 g, mobile phosphorus (according to Chirikov) – 17.1 mg/kg of soil, exchange potassium (according to Chirikov) – 17.8 mg/kg of dry soil, pH of salt extract – 6.5, hydrolytic acidity – 1.44 mg-eq/100 g, the sum of exchange bases is 35.2 mg-eq/100 g. The hydrothermal regime over the years of research was contrasting. According to the Vinnytsia meteorological station, during the growing season of 2019, 317.0 mm fell, which is 28.0 mm less than the annual average and 23.0 mm more than the amount of precipitation in 2020. The highest amount of precipitation (323 mm) was recorded in 2021. The air temperature during all the years of research (2019, 2020, and 2021) was higher by 0.8, 0.1, and 0.2°C, respectively, compared to the perennial average.

The three-factor field experiment was conducted according to the following scheme:

Factor A – varieties: Laperla – early, Granada – medium-early, Memphis – mid-season.

Factor B – fertiliser: 1. Control (without fertilisers); 2. 40 t/ha of semi-rotted manure under the predecessor + $K_{56}Mg_{16}S_{300}+P_{30}$ (background); 3. Background + $N_{30}R_{30}K_{30}$ (local application); 4. Background + $N_{45}P_{45}K_{45}$ (local application); background + $N_{60}P_{60}K_{60}$ (scattering application).

Factor C – tuber fraction (size according to the largest transverse diameter, mm and weight, g): 1. size <28 mm, weight 25-50 g; 2. size 28-60 mm, weight 51-80 g; 3. size 60 mm, weight 81-100 g.

Potatoes were grown according to the generally accepted technology for the Forest-Steppe zone. The potato planting rate was: for the fraction of <28 mm –

2-2.15 t/ha, 28-60 mm – 3.6-3.75 t/ha, 60 mm – 5.4-5.55 t/ha. Basic seed material of potato varieties was used for the study. The predecessor was winter wheat. Semi-rotted manure was applied under the predecessor – 40 t/ha. Potassium magnesium ($K_{28}Mg_8S_{15}$) and superphosphate (P_{30}) were applied for the main potato cultivation. During planting, Azofoska ($N_{15}R_{15}K_{15}$) was applied either in a scattering technique for pre-sowing cultivation. During the growing season of potatoes, phenological observations of plant growth and development were conducted and the following phases were visually noted: germination, budding, flowering, and tops dying off (Kutsenko *et al.*, 2002). Individual potato productivity was determined according to generally accepted methods (Bondarchuk & Koltunov, 2019). The fractional composition of seed potato tubers was determined by selection from all variants of the experimental sites in the first and third repetition. The mass of one sample was 10 kg. Potato tubers in the selected samples were divided into three fractions according to the largest transverse diameter: less than 28 mm, 28-60 mm, and over 60 mm. The number of tubers of each fraction was counted, weighed, and determined as a percentage of the total number. Accounting of tuber yield and determination of potato seed productivity was conducted in accordance with generally accepted methods (Kucenko *et al.*, 2002). The obtained results were processed according to modern statistical methods using computer programmes Excel and Statistica 6.0.

RESULTS AND DISCUSSION

It is possible to ensure the full realisation of the genetic potential of potato varieties by creating optimal conditions for the growth and development of plants in accordance with their needs during the growing season. Individual potato productivity depending on fertiliser, the fraction of planting tubers, and varietal characteristics is shown in Table 1.

With an increase in the fraction of planting tubers, the number of stems in the Laperla variety increased from 3.1 to 3.6 pcs./bush, the number of tubers – from 6.1 to 6.5 pcs./bush, and the mass of tubers – from 358.1 to 391.3 g/bush on the version without fertiliser. Application of 40 t/ha of semi-rotted manure under the precursor, potassium magnesium $K_{56}Mg_{16}S_{300}$, and superphosphate (P_{30}), basic potato cultivation, and an increase in the fraction of planting tubers contributes to an increase in the number of stems from 3.3 to 3.9 pcs./bush, the number of tubers – from 6.9 to 7.3 pcs./bush, and the mass of tubers – from 438.2 to 484 g/bush. An increase in individual potato productivity was observed against the background of phosphorus-potassium fertiliser and the aftereffect of semi-rotted manure and the local application of Azofoska ($N_{30}R_{30}K_{30}$). In this variant, the number of stems increases from 3.5 to 4.0 pcs./bush, the number of tubers – from 7.1 to 7.6 pcs./bush, and the mass of tubers from one bush – from 499.8 to 548.7 g.

Table 1. Individual potato productivity depending on fertiliser, seed tuber fraction, and variety, average for 2019-2021

Fertiliser (factor C)	Tuber fraction, mm (Factor B)	Quantity, pcs./bush		Mass of tubers, g/bush
		stems	tubers	
Laperla variety – factor A				
Control (without fertilisers)	1	3.1	6.1	358.1
	2	3.4	6.3	374.2
	3	3.6	6.5	391.3
40 t/ha+K ₅₆ Mg ₁₆ S ₃₀ +P ₃₀ (background)	1	3.3	6.9	438.2
	2	3.6	7.1	460.1
	3	3.9	7.3	484
Background +N ₃₀ R ₃₀ K ₅₀ (local application)	1	3.5	7.1	499.8
	2	3.8	7.3	520.5
	3	4.0	7.6	548.7
Background +N ₄₅ P ₄₅ K ₄₅ (local application)	1	3.7	7.5	545.3
	2	4.0	7.8	570.2
	3	4.3	8.1	595.4
Background +N ₆₀ P ₆₀ K ₆₀ (scattering application)	1	3.6	7.2	515.5
	2	3.9	7.5	543.0
	3	4.2	7.8	571.0
Variety Granada – factor A				
Control (without fertilisers)	1	3.5	6.5	464.8
	2	3.7	6.7	483.1
	3	3.9	7.0	510.3
40 t/ha of manure +K ₅₆ Mg ₁₆ S ₃₀ +P ₃₀ (background)	1	3.7	7.8	581.9
	2	3.9	8.0	600.8
	3	4.1	8.3	629.1
Background +N ₃₀ R ₃₀ K ₅₀ (local application)	1	3.9	8.6	665.6
	2	4.2	8.8	688.2
	3	4.5	9.1	718.9
Background +N ₄₅ P ₄₅ K ₄₅ (local application)	1	4.2	9.0	702.9
	2	4.5	9.2	725.9
	3	4.8	9.5	755.3
Background + N ₆₀ P ₆₀ K ₆₀ (scattering application)	1	4.1	8.8	864.6
	2	4.3	9.0	707.4
	3	4.4	9.3	735.6
Memphis – factor a variety				
Control (without fertilisers)	1	3.2	6.2	390
	2	3.4	6.4	405.1
	3	3.6	6.7	428.8
40 t/ha of manure +K ₅₆ Mg ₁₆ S ₃₀ +P ₃₀ (background)	1	3.3	8.1	552.4
	2	3.6	8.2	569.1
	3	3.9	8.5	596.7
Background + N ₃₀ R ₃₀ K ₅₀ (local application)	1	3.5	8.2	577.3
	2	3.8	8.5	606.1
	3	4.1	8.8	632.7
Background +N ₄₅ P ₄₅ K ₄₅ (local application)	1	3.7	8.5	609.5
	2	4.0	8.9	645.3
	3	4.3	9.2	675.3
Background + N ₆₀ P ₆₀ K ₆₀ (scattering application)	1	3.6	8.5	605.2
	2	3.8	8.7	625.5
	3	4.2	9.0	653.4

Source: compiled by the authors

The highest indicators of individual productivity of potato plants are formed with an increase in the fraction of planting tubers against the background of phosphorus-potassium fertiliser and the aftereffect of semi-rotted manure and the local application of Azofoska (N_{45}, R_{45}, K_{45}). For these options, the number of stems per bush increases from 3.7 to 4.3 pcs., the number of tubers per bush – from 7.5 to 8.1 pcs., and the weight of tubers per bush – from 545.3 to 595.4 g. Application of Azofoska by scattering under pre-sowing cultivation (N_{60}, R_{60}, K_{60}) against the background of phosphorus-potassium fertiliser and the aftereffect of semi-rotted manure does not provide an increase in individual productivity indicators compared to the previous version. Therefore, it is more effective for plants to use nutrients when applying fertilisers locally compared to applying them in a scattering technique.

The highest indicators of individual productivity among the varieties under study were observed in the Granada Variety at all variants of the experiment. Thus, in the control (without fertilisation), the number of stems was 3.5-5.9 pcs., the number of tubers under the bush – 6.5-7.0 pcs., and the mass of tubers from one bush – 464.8-510.3 g. Against the background of the aftereffect of semi-rotted manure and phosphorus-potassium fertiliser for the main cultivation of Granada potatoes, an increase in the number of stems from 3.7 to 4.1 pcs., the number of tubers from one bush and from 7.8 to 8.3 pcs., and the mass of tubers from the bush from 581.9 to 629.1 g was observed, depending on the fraction of planting tubers. Local application of Azofoska in rows (N_{30}, R_{30}, K_{30}) against the background of the aftereffect of semi-rotted manure and phosphorus-potassium fertiliser provided an improvement in the indicators of individual productivity of the Granada variety. Thus, in this variant, the number of stems is 3.9-4.5 pcs./bush, the number of tubers is 8.6-9.1 pcs./bush, and the weight of tubers is 665.6-718.9 g/bush, depending on the fraction of planting tubers. The highest indicators of individual productivity of the Granada variety were identified in the local application of Azofoska (N_{45}, R_{45}, K_{45}) against the background of the aftereffect of

manure and phosphorus-potassium fertiliser. Thus, the number of stems increases from 4.2 to 4.8 pcs./bush, the number of tubers – from 9.0 to 9.5 pcs./bush, and the weight of tubers – from 702.9 to 755.3 g/bush, depending on the fraction of planting tubers.

Application of Azofoska by scattering under pre-sowing cultivation (N_{60}, R_{60}, K_{60}) against the background of the aftereffect of semi-rotted manure and phosphorus-potassium fertiliser does not provide an increase in the number of stems and tubers in the bush, the mass of tubers from one Granada potato bush compared to the previous version. This indicates a smaller efficiency of applying fertilisers in a scattering technique compared to the local one. When applied locally, fertilisers are placed at a certain depth of the soil with a better moisture regime, where a zone with an increased concentration of nutrients is created. This contributes to the full use of nutrients by plants during the growing season.

The number of stems in the bush also has a substantial impact on the potato yield. In the process of potato growth and development, each stem becomes an independent plant that has its own root system, forms stolons and tubers. A strong direct correlation was established between the number of stems and the number of tubers in the bush ($r=0.78$), with a coefficient of determination of 61.0 %. Thus, an increase in the first indicator leads to an increase in the second (Figure 1). The yield of potato tubers depends on the productivity of each main stem, the number of stems in the bush, and the number of plants per area unit.

During three years of research, the structure of the potato yield was determined depending on the fertiliser, the fraction of planting tubers and varietal characteristics (Fig 2-4). As a result of the research, it was identified that with an increase in the fraction of planting tubers (from <28 to 60 mm) in the structure of the potato yield, the share of the tubers with the largest transverse diameter of 28-60 mm increases. Thus, in the Laperla Variety, the proportion of tubers in control (without fertilisers) with the largest transverse diameter of 28-60 mm is 42-51.1%, depending on the increase in the fraction of planting tubers (Fig. 2).




Figure 1. Correlation between the number of stems and the number of tubers in the bush

Source: compiled by the authors




Figure 2. Structure of the Laperla potato yield depending on the fraction of seed tubers and fertiliser (2019-2021), %
Note: 1. Control (without fertilisers); 2. 40 t/ha semi-rotted manure under the predecessor + $K_{56}Mg_{16}S_{30}+P_{30}$ (background); 3. Background + $N_{30}R_{30}K_{30}$ (local application); 4. Background + $N_{45}P_{45}K_{45}$ (local application); 5. Background + $N_{60}P_{60}K_{60}$ (scattering application).
Source: compiled by the authors

An increase in the fraction of planting tubers in the control caused a decrease in the proportion of tubers with the largest transverse diameter of >60 mm from 35.7 to 32.5%. This indicates that small planting tubers usually form one or two main stems with a small number of stolons and tubers. However, during the harvesting period, such potato plants usually form larger tubers. Notably, when planted with large seed tubers, more stems and tubers are formed, but smaller. Local application of Azofoska ($N_{45}R_{45}K_{45}$) against the background of the aftereffect of semi-rotted manure and phosphorus-potassium fertiliser provides the highest indicators (44.1-53.5%)

of the yield of potato tubers of the Laperla variety with the size of the largest transverse diameter of 28-60 mm. This indicator exceeds that on the control by 2.1-2.4 %. The yield of tubers of the Laperla variety with the largest transverse diameter of >60 mm in this variant is 35.1-37.4%, which is 1.7-2.6% more compared to the control.

The highest rates of tuber yield (44.9-54.6%) for the largest transverse diameter of 28-60 mm were provided by the medium-early Granada variety with local application of Azofoska ($N_{45}R_{45}K_{45}$) against the background of the aftereffect of semi-rotted manure and phosphorus-potassium fertiliser (Fig. 3).




Figure 3. Structure of the Granada potato yield depending on the seed tuber fraction and fertiliser (2019-2021), %
Note: 1. Control (without fertilisers); 2. 40 t/ha of semi-rotted manure under the predecessor + $K_{56}Mg_{16}S_{30}+P_{30}$ (background); 3. Background + $N_{30}R_{30}K_{30}$ (local application); 4. Background + $N_{45}P_{45}K_{45}$ (local application); 5. Background + $N_{60}P_{60}K_{60}$ (scattering application).
Source: compiled by the authors

Notably, the proportion of tubers of this fraction varied depending on the size of the tubers of the planting material. Thus, the yield share of tubers of the Granada variety with the largest transverse diameter of >60 mm in this variant is 34.9-38.0%. The lowest yield rates of tubers (10.5-17.1%) for the largest transverse diameter of <28 mm were formed by the Granada variety depending on the fraction of planting material, which is 4.3-4.9% less compared to the control. The best developed were plants from seed tubers of a >60 mm fraction. However, the yield

structure (excluding seed material) was maximal in the case of planting slightly smaller tubers (fraction of 28-60 mm), so along with the average seed fraction of tubers, it is advisable to plant healthy small tubers (<28 mm).

The highest rates of tuber yield (44.9-54.3%) for the largest transverse diameter of 28-60 mm were provided by the mid-season Memphis variety with local application of Azofoska ($N_{45}R_{45}K_{45}$) against the background of the aftereffect of semi-rotted manure and phosphorus-potassium fertiliser (Fig. 4).




Figure 4. Structure of the Memphis potato yield depending on the fraction of seed tubers and fertilizer (2019-2021), %
Note: 1. Control (without fertilisers); 2. 40 t/ha semi-rotted manure under the predecessor + $K_{56}Mg_{16}S_{30}+P_{30}$ (background); 3. Background + $N_{30}R_{30}K_{30}$ (local application); 4. Background + $N_{45}P_{45}K_{45}$ (local application); 5. Background + $N_{60}P_{60}K_{60}$ (scattering application)

Source: compiled by the authors

The yield share of Memphis tubers with the largest transverse diameter of >60 mm in this variant is 35.3-37.5%, which is 1.3-2.2% more compared to the control. It is known that the yield of potatoes depends

on the fraction of planting tubers. The features of the formation of yields of three potato varieties of different ripeness groups depending on fertilizer and seed tuber fraction were examined (Fig. 5)




Figure 5. Yield of potato varieties of different ripeness groups depending on fertilizer and fractions seed tubers (2019-2021), t/ha

Note: 1. Laperla Variety; 2. Granada Variety; 3. Memphis Variety

Source: compiled by the authors

On average, over three years, the highest yield of tubers (41.13 t/ha) was provided by the medium-early potato variety Granada in experimental plots where Azofoska ($N_{45}R_{45}K_{45}$) was locally applied against the background of the aftereffect semi-rotted manure and phosphorus-potassium fertiliser, planted with tubers with a fraction of >60 mm. The increase in tuber yield was 32.6% compared to the control (without fertilisers). The use of seed tubers with a fraction of >60 mm for planting Granada potatoes increases the yield by 1.5 t/ha compared to the fraction of planting tubers of 28-60 mm with the same method and background of fertilisation. Therefore, when noting yield gains from planting large tubers, it is necessary to mention the consumption of planting material. That is, the difference in the consumption of planting material between the best two variants of the Granada potato variety in

the experiment was 1.85 t/ha. Consequently, the increase in consumption of planting material was higher than the resulting increase in the yield of tubers.

A similar pattern was obtained in the early-maturing Laperla potato variety and the medium-maturing Memphis variety, where the yield was 32.5 and 36.7 t/ha, respectively, which is 34.2 and 36.5% more compared to the control (without fertilisers). The yield increase for planting these varieties with large tubers was 1.4 and 1.6 t/ha, respectively, and the difference in seed material consumption was 1.8 t/ha. That is, the seed material consumption on the variants for planting potatoes of Laperla and Memphis varieties with tubers with a fraction of >60 mm was also larger compared to the yield increases of these two varieties. Yield of seed potato varieties of different ripeness groups depending on fertiliser and the fraction of planting tubers (Fig. 6).




Figure 6. Yield of seed potato varieties of different ripeness groups depending on fertiliser and seed tuber fraction (2019-2021), t/ha

Note: 1. Laperla Variety; 2. Granada Variety; 3. Memphis Variety

Source: compiled by the authors

The highest yield (22.45 t/ha) of seed potatoes (fraction of 28-60 mm) was provided by the Granada variety in experimental plots where Azofoska ($N_{45}R_{45}K_{45}$) was locally applied against the background of the aftereffect of semi-rotted manure and phosphorus-potassium fertiliser, planted with tubers with a fraction of >60 mm. For local application of Azofoska ($N_{45}R_{45}K_{45}$) against the background of the aftereffect of semi-rotted manure and phosphorus-potassium fertiliser, planting tubers with a fraction of 28-60 mm, the yield was 21.2 t/ha. However, given the previously justified inexpediency of planting potatoes with tubers of the maximum size (>60 mm), due to the substantial consumption of planting material that was not overpowered by yield increases, it is advisable to use seed tubers with a fraction of 28-60 mm.

It is possible to ensure the formation of high tuber yields and increase resistance to adverse environmental conditions by providing potato plants with mineral nutrients adapted to their needs (Petropoulos *et*

al., 2020; Salikhov *et al.*, 2021; Naumann & Pawelzik, 2023). A study established that the application of mineral fertilisers affects the formation of the number of tubers and their mass (Mhango *et al.*, 2021). In the paper of P. Meise *et al.* (2019), the role of nitrogen in potato yield formation is highlighted. Sicilian researchers S. Lombardo *et al.* (2020) substantiated the application of nitrogen fertilisers (N_{140}) on typical soils of the region under study to obtain maximum yields of potato tubers with high consumer properties and increase the profits of producers. The dependence of the formation of individual potato productivity indicators on the application of organic and mineral fertilisers is confirmed by the obtained experimental data.

Similar studies by A. Podhaietskyi *et al.* (2019) confirm that the development of a large number of tubers in potato plants depends on the genetic characteristics of the variety or hybrid. The results are consistent with studies conducted in Ethiopia. It was identified that

planting potatoes with large seed tubers (46-55 mm) increases the number of stems and tubers per plant (Ebrahim *et al.*, 2018). S. Nasir and B. Akassa (2018) also reported an increase in the number of stems and tubers per plant when planted with large seed tubers. An increase in the number of stems in one potato bush under the influence of mineral fertilisers was also noted in the study by Y. Ilchuk & R. Ilchuk (2021). Thus, applying mineral fertilisers ($N_{60}P_{60}K_{90}$) increased the number of stems in the Shchedryk bush by 9.1 and 13.2% compared to the control, depending on the size of the planting tubers.

According to researchers (Podhaietsky *et al.*, 2019), one of the main factors in the productivity of potato varieties is the number of tubers formed in one plant. The dependence of potato yield on the number of stems and tubers in the bush is confirmed by the conducted research.

The results of this study are also consistent with those of D. AL-Taey *et al.*, (2008) on the positive impact of the use of organic and mineral fertilisers on improving the agrophysical condition of the soil, enhancing plant growth and development, and crop yield. According to U. Demirel (2023), to maximise the genetic potential of potato varieties during the growing season, it is necessary to create optimal conditions in accordance with the needs of plants. V. Gamajunova *et al.* (2021) confirmed that the selection of varieties and optimisation of plants with nutrients provides an increase in the yield of potato tubers by 64.7% compared to the control. M. Kravchuk *et al.* (2019) identified that using the organic-mineral fertiliser system (manure, 50 t/ha + straw, 3 t/ha + 30 kg/ha + green manure, 22.5 t/ha + $N_{45}P_{50}K_{60}$) increases the yield of potato tubers by 28.3-30.6% compared to the control. The data obtained is also consistent with the studies by Polishchuk (2021a, 2021b), which report the formation of the highest indicators of tuber yield in the Serpanok variety at different planting rates (50-60 thousand units/ha) on variants with the application of organic and mineral fertilisers. Optimisation of elements of crop cultivation technology should ensure not only an increase in their yield but also the profitability of production (Kotelnytska, 2020). Therefore, when noting yield gains from planting large tubers, it is necessary to mention the consumption of planting material.

According to researchers (Singh *et al.*, 2003), small seed tubers lead to reduced productivity, while large tubers become uneconomical due to their increased cost. The findings of this study are also consistent with the data from obtained by V. Kumar *et al.* (2021). Thus, they received the highest net profit from planting potatoes with medium-sized seed tubers compared with small and large ones. Therefore, they also recommend using medium-sized tubers (35-45 mm) to obtain high yields of high-quality seed material and increase profitability in Northwestern India.

It was identified that planting potatoes with medium (35-45 mm) and large (45-55 mm) seed tubers provided the formation of yields of small tubers (<35 mm) up to

7.4 and 8.3 t/ha, respectively, and medium-sized tubers (35-45 mm) up to 16.6 and 17.3 t/ha, respectively. The highest yield of tubers (20.2 t/ha) of large size (>45 mm) was obtained by planting small seed tubers (25-55 mm).

CONCLUSIONS

The formation of individual productivity indicators of potato varieties (the number of stems and tubers per bush, the mass of tubers per bush) depends on the method and background of fertilisation, varietal characteristics, and the fraction of seed tubers. Local application of Azofoska ($N_{45}R_{45}K_{45}$) against the background of the aftereffect of semi-rotted manure and phosphorus-potassium fertiliser provides an increase in the number of stems and tubers in the bush, the mass of tubers per bush, and the yield of potato varieties of different ripeness groups Laperla, Granada, and Memphis in Forest-Steppe conditions. The maximum indicators of the yield structure and the number of stems of Laperla, Granada, and Memphis varieties were observed in the experiment, where with an increase in the fraction of planting tubers against the background of phosphorus-potassium fertiliser and the action of semi-rotted manure, local application of Azofoska was conducted ($N_{45}R_{45}K_{45}$).

The highest yield rates of tubers of the Laperla, Granada, and Memphis varieties, the size of which with the largest transverse diameter of 28-60 mm, were obtained in the experiment, where locally applied Azofoska ($N_{45}R_{45}K_{45}$) against the background of the aftereffect of semi-rotted manure and phosphorus-potassium fertiliser for increasing the fraction of planting material. The yield of tubers in the varieties Laperla, Granada, and Memphis was 44.1-53.5%, 44.9-54.6%, 44.9-54.3% accordingly, the proportion of tubers with the largest transverse diameter of >60 mm varied from 37.4 to 35.1%, from 38.0 to 34.9%, and from 37.5 to 35.3%, respectively.

For rational use of seed material, potato varieties of different ripeness groups Laperla, Granada, and Memphis should be planted with tubers with a fraction of 28-60 mm. The inexpediency of using seed tubers with a fraction of >60 mm is explained by an increase in the consumption of planting material compared to the obtained yield increases. In addition, the yield structure (excluding seed material) was maximal when used for planting seed tubers with a fraction of 28-60 mm. Further research should focus on identifying the features of the formation of seed productivity of potatoes depending on the fertiliser and fraction of planting tubers due to the emergence of new varieties, subject to a change in the climate of the growing area.

ACKNOWLEDGEMENTS

None.

CONFLICT OF INTEREST

None.

REFERENCES

- [1] AL-Taey, D.K.A., Al-Naely, I.J.C., & Kshash, B.H. (2019). [A study on the effects of water quality, cultivars, organic and chemical fertilizers on potato \(*Solanum tuberosum* L.\) growth and yield to calculate the economic feasibility](#). *Bulgarian Journal of Agricultural Science*, 25(6), 1239-1245.
- [2] Biliavska, L., Biliavskiy, Y., Mazur, O., & Mazur, O. (2021). [Adaptability and breeding value of soybean varieties of Poltava breeding](#). *Bulgarian Journal of Agricultural Science*, 27(2), 312-322.
- [3] Bondarchuk, A.A., & Koltunov, V.A. (2019). *Potato growing: Methodology of the research case*. Vinnytsia: TVOTY.
- [4] Demirel, U. (2023). Environmental requirements of potato and abiotic stress factors. In M. E. Çalişkan, A. Bakhsh, & K. Jabran (Eds). *Potato Production Worldwide* (pp. 71-86). London: Academic Press. [doi: 10.1016/B978-0-12-822925-5.00011-6](#).
- [5] Ebrahim, S., Mohammed, H., & Ayalew, T. (2018). Effects of seed tuber size on growth and yield performance of potato (*Solanum tuberosum* L.) varieties under field conditions. *African Journal of Agricultural Research*, 13(39), 2077-2086. [doi: 10.5897/AJAR2018.13405](#).
- [6] Gamajunova, V., Khonenko, L., & Iskakova, O. (2021). Optimisation of nutrition of early-maturing potato varieties on drip irrigation in the South of Ukraine. *Scientific Horizons*, 24(8), 47-55. [doi: 10.48077/scihor.24\(8\).2021.47-55](#).
- [7] Ilchuk, Y., & Ilchuk, R. (2021). [Peculiarities of growth and development of early-rating varieties of potatoes depending on the area of father and size of the planting fraction of tubers](#). *Sciences of Europe*, 62, 3-10.
- [8] Iskakova, O., & Gamajunova, V. (2021). [Using biopreparations to optimize potato nutrition in the Southern Steppe of Ukraine](#). *AgroLife Scientific Journal*, 10(1), 109-115.
- [9] Kotelnitska, A., Tymoshchuk, T., Kravchuk, M., Sayuk, O., & Nevmerzhytska, O. (2021). [Mineral nutrition optimization as a factor affecting blue lupine crop productivity under conditions of global climate warming](#). *Romanian Agricultural Research*, 38, 223-230
- [10] Kotelnitska, G. (2020). The economic efficiency of the technology elements to grow blue lupine in the context of Polissia. *Scientific Horizons*, 1(86), 22-28. [doi: 10.33249/2663-2144-2020-86-1-22-28](#).
- [11] Kravchuk, M., Kropyvnytskyi, R., Andryash, V., Klymchuk, V., & Mysko, K. (2019). Change in soil agrophysical indicators and potato productivity in soil protection agrotechnologies. *Scientific Horizons*, 11(84), 61-68. [doi: 10.33249/2663-2144-2019-84-11-61-68](#).
- [12] Kumar, V., & Aulakh, C.S. (2022). Effect of planting geometry and potato seed tuber size on weeds and potato tuber yield. *Indian Journal of Weed Science*, 54, 291-295. [doi: 10.5958/0974-8164.2022.00052.1](#).
- [13] Kumar, V., Aulakh, C.S., & Kaur, J. (2021). Effect of planting geometry and potato seed tuber size on weeds and potato tuber yield. *Indian Journal of Agricultural Sciences*, 91(10), 108-111. [doi: 10.5958/0974-8164.2022.00052.1](#).
- [14] Kutsenko, V.S., Osypchuk, A.A., & Podhaietskyi, A.A. (2002). *Methodological recommendations for conducting research with potatoes*. Nemishaieva: Instytut kartopliarstva UAAN.
- [15] Kyrlyuk, V., Tymoshchuk, T., Kotelnitska, H., Barladiuha, V., & Dolid, D. (2020). Supplies of productive moisture and yielding capacity of crops rotation depending on the systems of basic tillage and fertilizing. *Scientific Horizons*, 7(92), 141-148. [doi: 10.33249/2663-2144-2020-92-7-141-148](#).
- [16] Lombardo, S., Pandino, G., & Mauromicale, G. (2020). Optimizing nitrogen fertilization to improve qualitative performances and physiological and yield responses of potato (*Solanum tuberosum* L.). *Agronomy*, 10, article number 352. [doi: 10.3390/agronomy10030352](#).
- [17] Meise, P., Seddig, S., Uptmoor, R., & Ordon, F. (2019). Assessment of yield and yield components of starch potato cultivars (*Solanum tuberosum* L.) under nitrogen deficiency and drought stress conditions. *Potato Research*, 62, 193-220. [doi: 10.1007/s11540-018-9407-y](#).
- [18] Mhango, J.K., Hartley, W., Harris, W.E., & Monaghan, J.M. (2021). Comparison of potato (*Solanum tuberosum* L.) tuber size distribution fitting methods and evaluation of the relationship between soil properties and estimated distribution parameters. *The Journal of Agricultural Science*, 159, 643-657. [doi: 10.1017/S0021859621000952](#).
- [19] Nasir, S., & Akassa, B. (2018). Review on effect of population density and tuber size on yield components and yield of potato (*Solanum tuberosum* L.). *African Journal of Plant Science*, 12(12), 319-23. [doi: 10.5897/AJPS2018.1701](#).
- [20] Naumann, M., & Pawelzik, E. (2023). Chapter13 - Nutrient management in potato. In M.E. Çalişkan, A. Bakhsh, & K. Jabran (Eds). *Potato Production Worldwide* (pp. 101-120), London: Academic Press. [doi: 10.1016/B978-0-12-822925-5.00018-9](#).

- [21] Petropoulos, S.A., Fernandes, Â., Polyzos, N., Antoniadis, V., Barros, L., C.F.R., & Ferreira, I. (2020). The impact of fertilization regime on the crop performance and chemical composition of potato (*Solanum tuberosum* L.) cultivated in central Greece. *Agronomy*, 10, article number 474. doi: [10.3390/agronomy10040474](https://doi.org/10.3390/agronomy10040474).
- [22] Podhaietskyi, A., Gnitetskyi, M., & Parchomenko, I. (2019). The ability to form potatoes of the progeny of interspecific and intraspecific potato hybrids. *Scientific Horizons*, 11(84), 69-76. doi: [10.33249/2663-2144-2019-84-11-69-76](https://doi.org/10.33249/2663-2144-2019-84-11-69-76).
- [23] Polishchuk, M. (2021a). The effect of fertilization system on the productivity of Serpanok – early mature variety of potatoes under the conditions of Right-Bank Forest-Steppe of Ukraine. *The Scientific Heritage*, 3(68), 18-28. doi: [10.24412/9215-0365-2021-68-3-18-28](https://doi.org/10.24412/9215-0365-2021-68-3-18-28).
- [24] Polishchuk, M. (2021b). The effect of sowing dates and nitrogen foliar fertilization on spring barley productivity under conditions of Right-Bank Forest-Steppe. *Agriculture and Forestry*, 23, 203-215. doi: [10.37128/2707-5826-2021-4-17](https://doi.org/10.37128/2707-5826-2021-4-17).
- [25] Sadawarti, M.J., Singh, S.P., Singh, R.K., Katare, S., & Samadhiya, R.K. (2021). Agro -techniques for production of seed size tubers in conventional seed potato production system – a review. *International Journal of Bio-resource and Stress Management*, 12(3), 238-246. doi: [10.23910/1.2021.2272](https://doi.org/10.23910/1.2021.2272).
- [26] Salikhov, T., Elubaev, S., Tynykulov, M., Kapbassova, G., & Makhmutova, A. (2021). The effect of the timing of manure application in combination with mineral fertilizers and planting density on the weediness of potato plantings. *Scientific Horizons*, 24(7), 46-52. doi: [10.48077/scihor.24\(7\).2021.46-52](https://doi.org/10.48077/scihor.24(7).2021.46-52).
- [27] Singh, A.K., Yada, V.K., Mir, M.S., & Khan, Z.H. (2003). [Standardization of seed size and spacing for improving yield of potato \(*Solanum tuberosum* L.\) under cold arid](#). *Environment and Ecology*, 21, 639-41.

Формування урожайності насінневої картоплі залежно від елементів технології вирощування

Ганна Володимирівна Миронова

Аспірантка

Вінницький національний аграрний університет
21008, вул. Сонячна, 3, м. Вінниця, Україна
<https://orcid.org/0000-0001-8401-6138>

Тетяна Миколаївна Тимошук

Кандидат сільськогосподарських наук, доцент
Поліський національний університет
10008, б-р Старий, 7, м. Житомир, Україна
<https://orcid.org/0000-0001-8980-7334>

Оксана Володимирівна Волошина

Кандидат філологічних наук, доцент
Вінницький національний аграрний університет
21008, вул. Сонячна, 3, м. Вінниця, Україна
<https://orcid.org/0000-0002-7679-9555>

Олена Василівна Мазур

Кандидат сільськогосподарських наук, доцент
Вінницький національний аграрний університет
21008, вул. Сонячна, 3, м. Вінниця, Україна
<https://orcid.org/0000-0001-6928-1865>

Олександр Васильович Мазур

Кандидат сільськогосподарських наук, доцент
Вінницький національний аграрний університет
21008, вул. Сонячна, 3, м. Вінниця, Україна
<https://orcid.org/0000-0002-2237-5116>

Анотація. Реалізувати максимальний генетичний потенціал сучасних сортів картоплі можна за рахунок удосконалення елементів агротехнології, зокрема оптимізації живлення рослин, що зумовлює актуальність цього дослідження. Метою роботи було обґрунтувати особливості формування продуктивності картоплі залежно від елементів технології вирощування в умовах Лісостепу. Польові дослідження проводили впродовж 2019-2021 рр. на чорноземних ґрунтах. Досліджено особливості росту і розвитку сортів картоплі різних груп стиглості Лаперла, Гранада і Мемфіс в умовах Лісостепу. Встановлено, що урожайність та вихід насінневих бульб картоплі досліджуваних сортів змінюється залежно від дози і способу внесення добрив, фракції садивних бульб та сортових особливостей. Найвищі показники індивідуальної продуктивності рослин картоплі сортів Лаперла, Гранада і Мемфіс сформовано за локального внесення в рядки $N_{45}P_{45}K_{45}$ на фоні дії напівперепрілого гною і фосфорно-калійного удобрення та використання насінневих бульб з фракцією >60 мм. У результаті дії досліджуваних факторів збільшується кількість стебел у кущі, кількість та маса бульб під одним кущем. Встановлено високої сили прямий кореляційний зв'язок між кількістю стебел у кущі і кількістю бульб ($r=0,78$), з коефіцієнтом детермінації 61,0%. Максимальну урожайність бульб картоплі досліджуваних сортів картоплі (29,7-41,1 т/га) отримали за локального внесення $N_{45}P_{45}K_{45}$ на фоні дії напівперепрілого гною і фосфорно-калійного удобрення залежно від розміру насінневих бульб за найбільшим поперечним діаметром. З'ясовано, що використання бульб фракцією >60 мм для садіння досліджуваних сортів картоплі є не раціональним, оскільки витрати насінневого матеріалу перевищують показники приросту урожайності. Визначено, що оптимальним для садіння сортів картоплі різних груп стиглості Лаперла, Гранада і Мемфіс бульб з фракцією 28-60 мм. Результати досліджень можуть бути використані для корегування елементів агротехнології насінневої картоплі, отримання якісного садивного матеріалу і введення прибутного агробізнесу

Ключові слова: удобрення; структура врожаю; індивідуальна продуктивність рослин; сорт; розмір бульб; фракція бульб
