

ISSN 2224-4980



**International
Journal of Ecosystems and Ecology Science
(IJEES)**

<https://doi.org/10.31407/ijeess>



Essays on Ecosystems and Environmental Research

Volume 13/2, 2023

February 2023 - June 2023

<http://ijeess.net/>

**International
Journal of Ecosystems and Ecology Science
(IJEES)**

<https://doi.org/10.31407/ijeess>





Clarivate Analytics

<https://rdmum.files.wordpress.com/2020/01/journal-list-emerging-sources-citation-index-web-of-science-update-on-23-jan-2020.pdf>

Volume 13, Issue 2, February 2023 - June 2023, (Serial Number 50)



Editor in Chief

Hysen Mankolli, Professor, Ecologist, Plainfield, Illinois, USA; Health and Environment Association, Tirana Albania.
Earth System Science Interdisciplinary Center (ESSIC), University of Maryland College Park, USA, Consultant Scientist.

Deputy-Editor-in-Chief

Sukru Dursun, Professor, Konya Technical University, Environmental Engineering Dep., Konya-Turkey.

Massimo Zuchetti, Professor, MIT, Massachusetts Institute of Technology, Cambridge MA, USA.

Cezar Kongoli, Professor, University of Maryland College Park, Earth System Science Interdisciplinary Center (ESSIC), USA.

Selim Dogan, Professor, Konya Technical University, Environmental Engineering Dep., Konya-Turkey.

International Scientific Editorial Board

Sukru Dursun, Professor, Konya Technical University, Environmental Engineering Dep., Konya, Turkey;

Massimo Zuchetti, Professor, MIT, Massachusetts Institute of Technology, Cambridge MA, USA;

Veselin Alexandrov, Professor, National Institute of Meteorology and Hydrology, Bulgaria;

Nicola Senesi, Professor, University of Bari, Bari, Italy;

Wander Michelle M, Professor, University of Illinois at Urbana-Champaign, Urbana, Urbana, IL 61801, USA;

Vladimir Pesic, Professor, University of Montenegro, Department of Biology, Podgorica Montenegro;

Ibraim Dincer, Professor, University of Ontario, Faculty of Engineering and Applied Science, Ontario, Canada;

Hysen Mankolli, Professor, Ecologist, Plainfield, Illinois, USA;

Lyudmyla Symochko, Prof. Assoc. Dr. Uzhhorod National University, Faculty of Biology, Uzhhorod, Ukraine;

Zacharoula S. Andreopoulou, Professor, Aristotle University of Thessaloniki, Thessaloniki, Greece;

Mariana Golumbeanu, PhD, National Institute for Marine Research and Development "Grigore Antipa", Constanta, Romania;

Antonis K. Kokkinakis, Professor, Aristotle University of Thessaloniki, Thessaloniki, Greece;

Olivia Cioboiu, PhD, The Oltenia Museum Craiova, Popa Șapcă str. No 8, Rumania;

Nasser Modirshahla, Professor, Islamic Azad University, Department of Applied Chemistry, Iran;

Nina Liogchii, Professor, Institut of Ecology and Geografy, Republic of Moldova;

Trajce Talevski, Professor, Hydrobiological Institute, Naum Ohidski 50, 6000 Ohrid - R.Macedonia;

Alexander V. Derunkov, Professor, Dep. of Entomology, National Museum of Natural History, Washington DC, USA;

Can Ozgur Colpan, Professor, Ryerson University, Canada;

Muhammad Ashraf, Professor, University of Agriculture, Faculty of Sciences, Faisalabad 38040, Pakistan;

Roberto Mancinelli, Professor, University of Tuscia, Viterbo, Italy;

Bahrive Gulgun, Professor, Ege University, Izmir, Turkey;

Insaf Mekki, PhD, National Research Institute for Rural Engineering, Water, and Forestry (I.N.R.G.R.E.F), Ariana, Tunisia;

Hany Gaber El Shaer, PhD, IUCN Centre for Mediterranean Cooperation / Global Marine, Egypt;

Ertugrul Esmeray, Ph.D, Karabuk University, Engineering Faculty, Environmental Engineering Department Turkey;

Violeta Vidaček-Hainš, PhD, Ass. Professor, University of Zagreb, Croatia;

Cezar Kongoli, Professor, University of Maryland College Park, Earth System Science Interdisciplinary Center (ESSIC), USA;

Justas Kažys, PhD, Assoc. Professor, Vilnius University, Department of Hydrology and Climatology, Vilnius –Lithuania;

Selim Dogan, PhD, Assoc. Professor Selcuk University, Environmental Engineering Department, Konya Turkey;

Hisham Mostafa Alidrisi, PhD, Assoc. Professor, King Abdulaziz University, Engineering College, Jeddah Saudi, Arabia;

Khalid A. A. Al-Ghamdi, PhD, Assoc. Professor, King Abdulaziz University, Engineering College, Jeddah Saudi, Arabia;

Osman Taylan, PhD, Assoc. Professor, King Abdulaziz University, Engineering College, Jeddah Saudi, Arabia;

Robert J. Kuligowski, PhD, University of Maryland College P, Earth System Science Interdisciplinary Center (ESSIC), USA;

Mariia M. Fedoriak, Professor, Chernivtsi National University, Dep. of Ecology and Biomonitoring, Chernivtsi, Ukraine;

Narayan Ramappa Birasal, Associate Professor, KLE Society's G H College, Zoology Department, , Karnataka state, India;

Rana P. Singh, Professor, Babasaheb Bhimrao Ambedkar University, Dep. of Environmental Science, U.P., India;

Hayati AKMAN, Dr., Selcuk University, Seed Department, 42430 Konya, Turkey;

Fatma ÇANKA KILIÇ, Professor, Kocaeli University, Technology Faculty, Energy Systems Engineering Dep., Kocaeli, Turkey;

Ferim Gashi, Dr., University of Pristina, Department of Geography, Faculty of Mathematical-Natural Sciences, Kosovo;

Altin Dorri, Professor Assoc., Polytechnic University of Tirana, Tirana, Albania;

Alexander P. Sizykh, Professor, Russian Academy of Sciences, Siberian Institute, Irkutsk, Lermonyova, Russia;

Zeynep EREN, Professor, Ataturk University, Environmental Engineering Department, Erzurum, Turkey;

Sheida Korjani, PhD, Department of Architecture, Tabriz Branch, Islamic Azad University, Tabriz, Iran;

Maan Maarroof, PhD, Assistant Professor, University of Mosul, College of Veterinary Medicine, Iraq;

Koula Doukani, Professor, University of Ibn Khaldoun, Zaaroura, Tiaret, Algeria;

Svetla Petkova Gateva, PhD, Institute of Biodiversity and Ecosystem Research, Bulgarian Acad. of Sciences, Sofia, Bulgaria;

Rrahim Sejdiu, Professor Assistant, PhD, University of Applied Sciences-Ferizaj, Kosovo;

Mouloud BOUHOUBOU, Dr., ENS Assia Djebar - Constantine, Algeria;

Dinh Tran Ngoc Huy, Dr., Banking University HCMC, Ho Chi Minh city Vietnam-International University of Japan, Japan;

Admir Jançe, Professor Assoc., European University of Tirana, Tirana, Albania;

Nataliia P. Kovalenko, Dr., Institute of Plant Physiology and Genetics of NAS of Ukraine, Kyiv, Ukraine;

Leila Soudani, Dr. University of Tiaret, Faculty of Nature and Life Sciences, Tiaret Algeria;

Publication Information:

International journal of ecosystems and ecology science (IJEES), ISSN 2224-4980, publishing original articles, reviews and short communications of high scientific standard on ecosystems and science ecology. By: Hysen Mankolli, Professor, Ecologist, Plainfield, Illinois, USA; Health and Environment Association, Tirana, Albania;
ISSN International Centre, *Bibliographic Data Section*, PARIS, FRANCE;
Bib - ID 68337; ISO standard 3297;
Key title: International journal of ecosystems and ecology science
Abbreviated key title: *Int. j. ecosyst. ecol. sci.*
ISSN 2224-4980; <http://ijeess.net/>
Frequency: four times a year. IJEES Electronic Journal Publication: Plainfield, Illinois, USA;
DOI prefix: 10.31407, <https://doi.org/10.31407/ijeess>



Clarivate Analytics

<https://rdmusm.files.wordpress.com/2020/01/journal-list-emerging-sources-citation-index-web-of-science-update-on-23-jan-2020.pdf>

Web of Science Core Collection,

Journal Citation Indicator (JCI): 2020: 0.05; 2019:0.05

<https://apps.clarivate.com/mjl-beta/search-results>

Emerging Sources Citation Index - JOURNAL LIST

<http://mjl.clarivate.com/cgi-bin/jrnlst/jlresults.cgi?PC=EX&Alpha=I>



ESCI Clarivate,

<https://rdmusm.files.wordpress.com/2020/01/journal-list-emerging-sources-citation-index-web-of-science-update-on-23-jan-2020.pdf>

UOI license: <http://u-o-i.org/1.01/ijeess>; <http://www.u-o-i.org/index.php/SearchLicense>



DPI Digital Library USA and assign unique digital no. DPI: 16.10047.IJEES

U.S. National Library of Medicine,

NLM ID: [101726509](https://pubmed.ncbi.nlm.nih.gov/101726509/) [Serial]; Other ID: (OCOLC)879805056;

<https://www.ncbi.nlm.nih.gov/nlmcatalog/101726509>

This journal has the status of an international journal.

Aims and Scope:

The goals of the International Journal of Ecosystems and Ecology Science (IJEES), ISSN 2224-4980, are to bring together researchers and scientists with interests in the quality of the ecosystems research results, theories, technologies, systems, tools, applications, the work in progress and experiences on ecosystems used. The main topics of interest are:

- ✓ Ecosystems
- ✓ Agro ecosystems
- ✓ Forest ecosystems
- ✓ Ecology
- ✓ Plant Ecology
- ✓ Animal Ecology
- ✓ Human Ecology
- ✓ Lakes and Rivers ecosystems
- ✓ Applied Biology
- ✓ Applied Ecology
- ✓ Applied Chemistry
- ✓ Biodiversity
- ✓ Energy
- ✓ Geology
- ✓ Bioaccumulation
- ✓ Desalination
- ✓ Water
- ✓ Soil
- ✓ Air pollution
- ✓ Climate Change
- ✓ Ecosystem restoration
- ✓ Environment toxicology
- ✓ Environment protection
- ✓ Environmental radioactivity
- ✓ Environmental legislation
- ✓ Environmental management
- ✓ Environmental education.
- ✓ Green technology
- ✓ Socioeconomic aspects in Ecosystems
- ✓ Agro tourism and National Park
- ✓ Health Care
- ✓ Food security
- ✓ Rights and Environmental Laws
- ✓ Environment Engineering
- ✓ Environment Architecture
- ✓ Eco philosophy Issue

Abstracted / Indexed / Impact Factor in:

International journal of ecosystems and ecology science (IJEES), ISSN 2224-4980,

IJEES: <http://ijeess.net/>

IJEES have an international status and has been indexed, abstracted, and have impact factors:

ISSN Center responsible of the record: CIEPS - ISSN

<https://portal.issn.org/resource/ISSN/2224-4980>

IJEES doi crossref

<https://doi.org/10.31407/ijeess>

IJEES, Volume 13/2 2023

[https://ijeess.net/journal-89-International-Journal-of-Ecosystems-and-Ecology-Science-\(IJEES\)-Volume-13-2,-2023.html](https://ijeess.net/journal-89-International-Journal-of-Ecosystems-and-Ecology-Science-(IJEES)-Volume-13-2,-2023.html)

Web of Science Master List

<https://rdmum.files.wordpress.com/2020/01/journal-list-emerging-sources-citation-index-web-of-science-update-on-23-jan-2020.pdf>

Journal Citation Report™ (JCR), Journal Citation Indicator (JCI):2021: **0.04**; 2020:**0.05**

<https://clarivate.com/webofsciencelibrary/solutions/journal-citation-reports/>

Emerging Sources Citation Index - JOURNAL LIST

<http://mjil.clarivate.com/cgi-bin/jrnlst/jlresults.cgi?PC=EX&Alpha=I>

Sciencegate cited

<https://www.sciencegate.app/source/315335/top-cited>

Scilit

<https://www.scilit.net/journal-articles/1892436>

CiteFactor: IF 2019-2020:1.40;

<https://www.citefactor.org/impact-factor/impact-factor-of-journal-International-Journal-of-Ecosystems-and-Ecology-Science-IJEES.php>

EBSCO's library products,USA

<http://www.ebscohost.com/for-publishers/for-publishers>

IARC: Journal Impact Factor (USA), IF=2017: **1.811**; IF=2016: **1.463**; =2015: **0.861**

http://iarconf.org/index.php/journal_information/show/44/0/2224-4980

Index Copernicus ICV= 2021: **100.00**; ICV= 2020: **100.00**; ICV= 2019: **100.00**;

<https://journals.indexcopernicus.com/search/journal/issue?issueId=197247&journalId=31407>

Academic Resource Index Research Bible:

<http://paper.researchbib.com/view/issn/2224-4980/10/2>

Scientific Journal Impact Factor Value 2020: **SJIF 2020 = 7.167**

<http://sjifactor.com/masterlist.php?area=all&country=al&ord=sjif>

Google scholar Cited: **h-index: 10**; **Citations 400**;

<https://scholar.google.com/citations?user=W89N8qgAAAAJ&hl=en>

MIAR

<http://miar.ub.edu/issn/2224-4980>

Directory of Research Journals Indexing (DRJI)

<http://olddrji.lbp.world/JournalProfile.aspx?jid=2224-4980>

WorldCat

https://www.worldcat.org/search?qt=affiliate_wc_org_all&ai=Directory_ashok.drji%252540gmail.com&fq=&q=2224-4980&wscbtn2w=Go

US. National Library of Medicine, NLM ID: [101726509](https://pubmed.ncbi.nlm.nih.gov/101726509/) [Serial]; Other ID: (OCOLC)879805056;

<https://www.ncbi.nlm.nih.gov/nlmcatalog/101726509>

Global Impact Factor (GIF) IF: 2015: **0.765**; 2014: **0.625**; 2013: **0.541**; IF= 2012: **0.425**;

<http://globalimpactfactor.com/journals-list/?snap=I>

CAS Chemical Abstracts Service – CAS database

<http://cassi.cas.org/search.jsp>

Journal Factor: ISSN 2224-4980

http://www.journalfactor.org/advance_search.php

Directory of Research Journals Indexing (DRJI)

<http://olddrji.lbp.world/justIncluded.aspx>

Environment-and-Nature/Biodiversity/Ecosystem.

<http://www.efita.org/Environment-and-Nature/Biodiversity/Ecosystem-Biodiversity/IJEES-details-28892.html>

NewJour : Electronic Journal&Newsletters

<http://gulib.georgetown.edu/newjour/nj2/index.html>

Environmental expert

<http://www.environmental-expert.com/magazines/?keyword=ecosystem+science>

Currently Indexing at Pro Quest

http://www.proquest.com/en-US/catalogs/databases/detail/periodicals_index.shtml

CABI

<http://www.cabi.org/default.aspx?page=1016&site=170&pid=40&xslttab=2&newtitlesonly=1&search=>

Manuscripts and correspondence are invited for publication

Dear Professors, PhD, researchers and scholars of the field of ecosystems and related areas to, such as environmental engineering, agriculture, bio-chemistry, soil, water and air pollution, geology, forestry, urban ecology, resource economics, etc., at the moment, I declare the sending of scientific journal articles to International Journal of Ecosystems and Ecology Science (IJEES) is open. Sending of items can only be done electronically, to the following address: e-mail: editorijeess@gmail.com; 2011ijeese@gmail.com; mjedisishendeti@yahoo.com; The confirmation of receiving these items will be announced within the period of one to three weeks after sending. Whether the article is accepted or not, will be announced within a period of two to three weeks. Only original articles will be accepted, validated, and not to have or be published in other journals. Publication lasts from two to six months. All Rights Reserved Copyright by © IJEES (the first or correspondent author have responsible for the accuracy of the article). Submission guidelines and Web Submission System are available at: <https://www.ijeess.net/page-6-Guide-for-authors.html>;
Note: If the paper is published in another journal or if it includes false or inaccurate claims, it will be rejected or removed automatically as that constitutes a violation. Zero tolerance!

Editorial Office:

IJEES Electronic Journal Publication: Plainfield, Illinois, USA; E-mail: editorijeess@gmail.com;
Health and Environment Association, Tirana, Albania; E-mail: mjedisishendeti@yahoo.com;

**International
Journal of Ecosystems and Ecology Science
(IJEES)**

DOI: <https://doi.org/10.31407/ijeess>

Volume 13/2, 2023

DOI: <https://doi.org/10.31407/ijeess13.2>

Table of contents:

Oleksandr Tkachuk*, Nadiya Viter, Snizhana Pankova, Olga Titarenko, Lyudmila Yakovets, THE CURRENT ENVIRONMENTAL STATE OF THE FIELD PROTECTIVE FOREST BELTS OF THE FOREST STEPPE OF UKRAINE, page 1-6;

DOI: <https://doi.org/10.31407/ijeess13.201>

Mohammed Balubaid¹, Waqar Ahmad Gulzar¹, Hani Aburas¹, Osman Taylan¹, Abdulaziz S. Alkabaa¹, Omer A. Bafail¹, Anas A. Makki², Ammar Y. Alqahtani¹, Hassan M. Alidrisi¹, Basil Omar Al-Sasi¹, Shoukath Ali Karuvatt³, Hisham Alidrisi^{1*}, MONITORING THE PERFORMANCE OF AGRICULTURAL AND FOOD SECTOR COMPANIES USING DEA, page 9-24;

DOI: <https://doi.org/10.31407/ijeess13.202>

DOUKANI Koula^{12*}, BOUHENNI Hasna¹, BOUKIRAT Dyhia³, HADJARAB Mohamed Hocine¹, MOKHTAR Asma¹, ZAHI Racha¹, EPIDEMIOLOGICAL STUDY OF GASTRODUODENAL DISEASES (*HELICOBACTER PYLORI*) IN THE REGION OF TIARET, ALGERIA, page 25-32;

DOI: <https://doi.org/10.31407/ijeess13.203>

Eugena Tomini^{1*}, Adela Vasili¹, Elona Kureta¹, Artan Simaku¹, Silvia Bino¹, COVID-19: A PANDEMIC EXPERIENCE FOR THE COOPERATION OF THE SCIENTIFIC COMMUNITY WORLDWIDE, page 33-36;

DOI: <https://doi.org/10.31407/ijeess13.204>

M Balubaid1, W A Gulzar1, H Aburas1, O Taylan1, A S Alkabaa1, O A Bafail1, A A Makki2, A Y Alqahtani1, H M Alidrisi1, B O Al-Sasi1, Sh A Karuvatt3, H Alidrisi1*, MONITORING THE PERFORMANCE OF AGRICUL. AND FOOD SECTOR COMPANIES USING DEA, page 9-24; -

DOUKANI Koula12*, BOUHENNI Hasna1, BOUKIRAT Dyhia3, HADJARAB Mohamed Hocine1, MOKHTAR Asma1, ZAHI Racha1, EPIDEMIOLOGICAL STUDY OF GASTRODUODENAL DISEASES (HELICOBACTER PYLORI) IN THE REGION OF TIARET, ALGERIA, page 25-32; -

Eugena Tomini1*, Adela Vasili1, Elona Kurefa1, Artan Simaku1, Silvia Bino1, COVID-19: A PANDEMIC EXPERIENCE FOR THE COOPERATION OF THE SCIENTIFIC COMMUNITY WORLDWIDE, page 33-36; -

Fatma Köprülü1*, Osama Abduljalil Mohammad Hamad2*, Aşkın Kiraz3*, PERCEPTIONS OF LIBYANS TOWARDS CLIMATE CHANGE, page 37-44; -

Nevila Bushati1,2*, Anila Neziri2, Arifind Mustafa3, PHYSICO-CHEMICAL AND BACTERIOLOGICAL EVALUATION FROM GROUNDWATERS OF PUKA CITY, page 45-50; -

Inna Honcharuk, Oleksandr Tkachuk*, Olexandr Mazur, Ruslan Kravets, Olena Mazur, Oleksiy Alieksieiev, Zabarna Tetyana, Lina Bronnikova, MANIFESTATION OF ECOLOGICAL-ADAPTIVE PROPERTIES OF SOYBEAN VARIETIES DEPENDING ON SOIL-CLIMATIC CONDITIONS, page 51-60; -

Mohammed Alamoudi1*, SAFETY CLIMATE DIMENSIONS AS PREDICTORS FOR WORKER'S SATISFACTION REGARDING SAFETY LEVEL, page 61-70; -

S Razanov1,2*, O Koruniak3, A Dydiv1, T Holubieva4, L Symochko5,6, V Balkovskiy1, O Alekseev2, O Vradii2, H Ohorodnichuk2, M Polishchuk2, O Kolisnyk2, O Mazur2, O Mazur2, RADIOPROTECTIVE AND SORPTION PROPERTIES OF BEE SPINE, page 71-76; -

Furkan TÜRK1,2, Hayriye Nur NAYAN2, Selim DOĞAN3*, Ülkü Sultan KESKİN2, THE CARBON COST OF THE BUILDINGS DESTROYED IN YALOVA IN THE 1999 MARMARA EARTHQUAKE, page 85-90; -

V Mazur1, H Pantsyreva1*, O Ovcharuk2, K Mazur1, R Myalkovsky3, O Tkach3, S Verholiuk1, FEATURES OF THE FUNCTIONING OF THE ASSIMILATION APPARATUS OF LUPINUS ALBUS DEPENDING ON THE USE OF ECO-BIOLOGICAL PREPARATIONS, page 77-84; -

Muhamet Kadrija1*, Sabrije Kelmendi-Bardhaj2, Besarta Pelaj3, Haxhi Kamberi4*, NURSING CARE FOR PATIENTS AT THE END LIFE, page 109-124; -

MUNTING DIVINE TII1*, TCHOBSALA1, LOCAL EXPLOITATION AND CHARACTERIZATION OF BALANITES AEGYPTIACA (L.) DEL PARKS IN THE SUDANO-SAHELIAN ZONE OF CAMEROON, FAR-NORTH, CAMEROON, page 125-138; -

Alla Levishko1, Iryna Gumeniuk1*, Viktoriya Tsvigun1, Svitlana Mazur1, Yeuheniia Tkach1, Olena Demyanyuk1, Tetiana Kovalenko2, THE INFLUENCE OF FIRE CULTIVATION ON THE SOIL MICROORGANISMS, page 91-96; -

Oleg Tkach1, Viktor Dubik1, Oleh Ovcharuk2, Lyudmila Mikhaylova1, Hanna Pantsyreva3*, Dariia Vilchynska1, Sergii Slobodian1, Oleg Gorbovy1, TECHNOLOGICAL CHARACTERISTICS AND POTENTIAL OF LANDFILL BIOGAS FOR ELECTRICITY GENERATION, page 97-108; -

Ana T. Misoska1, Snezhana Hristova1*, Elena Bundaleska1, Sanja Georgieva1, HOW DO EMPLOYEES' ATTITUDES TOWARDS "GREEN BEHAVIOR" RELATE TO ENVIRONMENTAL CULTURE.. page 139-152; -

FEATURES OF THE FUNCTIONING OF THE ASSIMILATION APPARATUS OF LUPINUS ALBUS DEPENDING ON THE USE OF ECO-BIOLOGICAL PREPARATIONS

Victor Mazur¹, Hanna Pantsyreva^{1*}, Oleh Ovcharuk², Kateryna Mazur¹,
Ruslan Myalkovsky³, Oleg Tkach³, Serhii Verholiuk¹

¹*Vinnitsia National Agrarian University, Vinnitsia, Ukraine;*

²*National University of Life and Environmental Sciences, Ukraine;*

³*State Agrarian and Engineering University in Podilya, Ukraine;*

*Corresponding Author Hanna Pantsyreva, email: apantsyreva@ukr.net;

Received January 2023; Accepted February 2023; Published March 2023;

DOI: <https://doi.org/10.31407/ijeess13.210>

ABSTRACT

The high efficiency and justified necessity of using biopreparations for pre-sowing seed treatment and foliar fertilization has been proven. The results of studies on the effectiveness of the application of pre-sowing treatment and foliar feeding of white lupine plants in the conditions of the right-bank forest-steppe of Ukraine are presented. The optimal leaf surface area and photosynthetic potential were determined. A positive effect of pre-sowing seed treatment with the bacterial preparation Rhyzogumin and the growth stimulator Emistym C in combination with two foliar feedings Emistym C on the content of chlorophyll in the leaves of white lupine varieties Veresnevyi and Makarivskyi was established. Pre-sowing treatment of white lupine seeds of the Veresnevyi and Makarivskyi varieties in combination with foliar feeding contributed to the increase of the leaf surface area and the formation of the photosynthetic apparatus of plants. The effect of the investigated biological preparations on the formation of the assimilation surface and the synthesis of chlorophyll in the leaves of white lupine in the conditions of the right-bank forest-steppe of Ukraine has been proven. It was determined that the greatest stimulatory effect was obtained in the variant where the pre-sowing treatment of seeds involved the interaction of the biological preparations Rhyzogumin and Emistym C in combination with two foliar feedings of Emistym C. The three-year research results also indicate a significant impact of the researched technological methods of cultivation on grain yield.

Keywords: white lupine, variety, pre-sowing seed treatment, foliar feeding, assimilation apparatus, leaf area, photosynthetic potential, chlorophyll.

INTRODUCTION

Lupine is a culture of universal possibilities. Along with providing valuable feed raw materials, lupins are of great importance in increasing soil fertility, phytomelioration, improving the phytosanitary condition of agrocenoses, and reducing energy consumption in crop production (Mazur et al., 2019; Pancy'reva, 2016; Kaletnik & Lutkovska, 2020). White lupine, compared to other types of fodder lupine, is characterized by fast growth rates, early maturity and high fodder and grain productivity. Cultivation of lupine provides an increase in soil fertility, replenishment of

the soil nitrogen balance due to its biological fixation, and contributes to the strengthening of the financial condition of farmers. In terms of chemical composition and nutritional value, lupine protein is closest to animal protein. The high content of valuable protein in the plant and a complex of other economically valuable features make lupine an irreplaceable fodder crop. Currently, in the countries where white lupine is grown, numerous scientific studies are being conducted, which are aimed at a detailed study of the chemical composition of this plant. At the same time, the results of studies of white lupine must be compared with soybeans in order to prove their competitiveness and study the possibility of an alternative replacement. In terms of seed quality, lupine is not inferior to this crop, so it is called the second northern soybean (Didur et al., 2020). Formation of grain productivity and fodder value of agricultural crops largely depends on the intensity of photosynthesis, synthesis and transport of metabolites in leaves. Therefore, it is possible to increase the realization of the potential of plants due to the activation of these processes, in particular, the process of photosynthesis. Formation of productivity as a result of photosynthetic activity of plants in crops is determined by the functioning of the assimilation apparatus (Zhu et al., 2019; Ahmadi et al., 2010). Scientifically based bases of technologies for growing leguminous crops, including white lupine, determining the accumulation of chlorophyll in plant leaves is important, since their content affects the intensity of photosynthesis and other physiological processes. The research is aimed at establishing the peculiarities of the functioning of the photosynthetic apparatus, the peculiarities of the formation of the assimilation apparatus in the processes of growth and development of plants are of primary importance when assessing the impact of technological techniques on grain productivity and feed value of plants. In this regard, conducting relevant research is important in modern agricultural production

MATERIALS AND METHODS

Field experiments were conducted in the conditions of the research farm «Agronomichne» of the Vinnytsia National Agrarian University in 2018-2020. The experimental sites are located in the right-bank forest-steppe zone of Ukraine. The soil cover is represented by gray forest soils with a humus content in the soil layer of 0-30 cm at the level of 2.09. The climate in the research area is temperate-continental with an annual precipitation of 418 mm and average monthly temperatures during the growing season of chickpeas in the range of 15.5-22.8 °C. During the years of the experimental experiments, the growing season was characterized by higher (by 0.9-4.3 °C) temperatures and a deficit of precipitation compared to the long-term norm, which was 56 mm less.

The purpose of the research is to establish the specifics of the formation of the assimilation apparatus by plants of white lupine (*Lupinus albus* L.) depending on biological preparations. The material for the research was white lupine (*Lupinus albus* L.) Veresnevyi and Makarivskiyi varieties.

The technology of growing white lupine varieties is generally accepted for the forest-steppe zone of Ukraine and included pre-sowing treatment of seeds with the bacterial preparation Rhyzogumin in combination with the growth stimulator Emistym C and foliar feeding Emistym C. On the day of sowing, white lupine seeds were treated with the bacterial preparation Rhyzogumin (600 g per hectare rate of seeds) and growth stimulator Emistim C (10 ml per 1 ton of seeds) using PKS-20 Super. In foliar top dressing, the growth stimulator Emistim C was used at a rate of 15 ml/ha. The first foliar top dressing of Emistym C was carried out in the budding phase, the second - in the phase of the beginning of pouring seeds. The area of the accounting plot is 25 m². Repetition - five times. Placement of options is systematic in two tiers.

Evaluation of the photosynthetic activity of plants was carried out according to the following methods: the area of the leaf surface was measured by the «cuts» method, the photosynthetic potential was determined according to the method of A. A. Nychiporovich (1996); the amount of chlorophyll was determined by the method of alcohol weighing on a conditioned electrophotocolorimeter (KFK-2) (Nychyporovych et al, 1961).

RESULTS AND DISCUSSION

Chlorophyll content in leaves is an indicator characterizing the potential level of photosynthetic productivity of plants. The informativeness of this parameter is determined by a set of features that do not allow using the leaf surface area as an absolute indicator of their photosynthetic productivity. First of all, it is the thickness of the leaf

plate and the total concentration of pigments in the cells. The last indicator can vary significantly depending on the species and even the variety. As a rule, species that spread from south to north are characterized by an increase in leaf surface area with a decrease in total chlorophyll concentration. Conversely, in species whose range expansion goes in the opposite direction – from north to south – a decrease in the area of individual leaves is observed while maintaining or increasing the concentration pigments (Chynchyk, 2013; Langewisch, 2017).

Domestic and foreign authors indicate that the biological harvest depends on the content of pigments, primarily chlorophylls in the assimilating organs of plants, time and intensity of their work. The content of chlorophyll in leaves affects the intensity of photosynthesis, the accumulation of dry matter, and ultimately their productivity. The need for research in this direction is due to the fact that the total mass of green pigment and its concentration in the mesophyll of the leaf, together with the dimensions of the assimilation surface, are considered as the basis of the potential of the photosynthetic activity of the plant organism as a whole (Vdovenko, 2018; Kantolic, 2013).

The difference in chlorophyll content, as a rule, is an indicator of the level of compliance with vegetation conditions and varies depending on the genotype of the variety. Increasing the yield of agricultural crops depends both on factors affecting photosynthesis and on the complex of physiological processes associated with it (water exchange, nutrition, growth). The formation of a well-developed photosynthetic apparatus, optimal in terms of volume, dynamics and intensity of functioning, is the key to the creation of organic matter, biological and marketable crops (Mazur et al., 2021; Kaletnik et al., 2020; Puyu et al., 2021).

Research has established that for the formation of a crop as a photosynthetic system, many factors should be taken into account, among which the variety, its ecology and biology, and technological methods of cultivation are of great importance (Rai et al., 2017; Bandura et al., 2019).

During the years of research, the dependence between the process of formation of the leaf surface of lupine and the elements of growing technology was observed. The results of the conducted research showed that the choice of varieties that are not damaged by pests and diseases, the phases of vegetation and technological methods were of great importance for the functioning of the leaf surface.

It was noted that the area of the leaf surface of white lupine increases gradually, reaching the maximum value in the phase of the beginning of seed pouring. After this phase, a decrease in leaf area was observed, which is determined by the biological characteristics of the culture. Thus, the redistribution and increased outflow of plastic substances from vegetative organs to seeds causes the death of leaves during the ripening of white lupine plants. A similar trend of the effect of the researched technological methods on the leaf surface area was also found in the plots of the Makarivskiy white lupine variety. However, the indicators of the dynamics of the leaf surface area were somewhat lower than in the Veresnevyi variety. This is explained by a genetically determined feature.

The formation of the size of the leaf area in different phases of growth and development of white lupine was influenced by the pre-sowing treatment of seeds with nodule bacteria and a growth stimulator in combination with foliar fertilization (Fig. 1).




Figure 1. Dynamics of the leaf surface area of white Lupine plants of the Veresnevyi variety depending on biological preparations, thousand m²/ha (2018-2020)

Thus, in the phases of branching and budding, the influence of the investigated technological techniques on the parameters of the leaf surface area was insignificant. On the plots of the Veresnevyi variety, during budding, the leaf area index, depending on the pre-sowing treatment of seeds and foliar fertilization, was in the range of 14.2-15.7 thousand m²/ha, and during the period of full flowering - from 22.3 to 27.6 thousand m²/ha. The formation of the photosynthetic surface of the crop is influenced by both biotic and abiotic factors. Optimizing thermal, water, air, and nutrient regimes is of great importance for the productive work of sowing as a photosynthetic system.

Photosynthesis is the main complex and multi-stage process of plant nutrition, therefore, other processes are effective only to the extent that they improve and stimulate photosynthetic activity and create conditions for the synthesis of assimilants and their best use for the processes of growth, development and crop formation. Organic substances formed in the process of photosynthesis make up about 95% of the dry mass of plants. Therefore, formation by the process of photosynthesis, increasing its productivity is one of the effective methods of influencing the productivity of plants, and for agricultural crops it is an important means of increasing the yield (Mazur et al., 2019; Honcharuk et al., 2020; Kaletnik, 2010; Mazur et al., 2020; Didur et al., 2020; Bulgakov et al., 2014). The records conducted during the research period showed that the pre-sowing treatment of lupine seeds with the bacterial preparation Rhizogumin in combination with the growth stimulator Emistym C with two foliar feedings Emistym C has a positive effect on the formation of the photosynthetic apparatus of white lupine plants (Table 1).

Table 1. The formation of the photosynthetic potential of white lupine varieties depends from biological preparations, million m²/ha (average for 2018-2020).

Factors			Vegetation periods of plants			
variety	pre-sowing seed processing	foliar feeding*	full stairs-budding	fulladder-fullbloom	full stairs-start of pouring	full stairs-physiological maturity
Veresnevyi	Without pre-sowing seed treatment**	without supplements**	0,321	0,601	0,989	1,505
		one feeding	0,321	0,606	1,005	1,529
		two feedings	0,321	0,606	1,006	1,559
	Rhizohumin	without supplements	0,326	0,616	1,050	1,588
		one feeding	0,326	0,622	1,075	1,638
		two feedings	0,336	0,622	1,076	1,689
	Emistym C	without supplements	0,337	0,637	1,125	1,766
		one feeding	0,337	0,648	1,150	1,819
		two feedings	0,337	0,648	1,151	1,860
	Rhizohumin + Emystim C	without supplements	0,354	0,675	1,125	1,941
		one feeding	0,354	0,689	1,260	1,982
		two feedings	0,354	0,689	1,262	2,061
Makarovskyi	Without pre-sowing seed treatment	without supplements	0,281	0,495	0,861	1,301
		one feeding	0,281	0,501	0,878	1,333
		two feedings	0,281	0,501	0,879	1,351
	Rhizohumin	without supplements	0,290	0,516	0,921	1,389
		one feeding	0,290	0,525	0,939	1,420
		two feedings	0,290	0,525	0,940	1,452
	Emistym C	without supplements	0,301	0,552	1,009	1,522
		one feeding	0,301	0,566	1,036	1,577
		two feedings	0,301	0,566	1,037	1,615
	Rhizohumin + Emystim C	without supplements	0,316	0,574	1,073	1,650
		one feeding	0,316	0,589	1,093	1,678
		two feedings	0,316	0,589	1,095	1,720

Notes: * – Emistym C; ** – CONTROL.

However, the effectiveness of the application of pre-sowing treatment of seeds and foliar fertilization is noticeable in the phase of physiological maturity. Thus, the highest indicators of photosynthetic potential of white lupine varieties Veresnevyi and Makarivskyi were observed in the period of full germination - physiological maturity in the variants with the use of a bacterial drug and a growth stimulator in the pre-sowing treatment of seeds in combination with two foliar feedings and were, respectively, 2.061 and 1.720. These indicators of white lupine plants of Veresnevyi and Makarivskyi varieties exceeded the control variant by 27.0% and 21.5%, respectively. The presence of a positive effect of pre-sowing seed treatment and foliar fertilization on the content of chlorophyll in white lupine leaves was established (Table 2).

Table 2. Chlorophyll content of white lupine varieties depending on biological preparations (average for 2018-2020)

Factors			Chlorophyll content in leaves, mg/g raw mass	Chlorophyll content in leaves, mg/m ²
variety	pre-sowing seed processing	foliar feeding*		
Veresnevyi	Without pre-sowing seed treatment**	without supplements**	2,03	2101,28
		one feeding	2,03	2101,28
		two feedings	2,07	2560,29
	Rhizohumin	without supplements	2,16	2699,44
		one feeding	2,23	2707,68
		two feedings	2,33	3246,51
	Emistym C	without supplements	2,16	2699,44
		one feeding	2,23	2707,68
		two feedings	2,48	3679,94
	Rhizohumin + Emistym C	without supplements	2,48	3679,94
		one feeding	2,69	4083,31
		two feedings	2,87	4802,12
Makarivskyi	Without pre-sowing seed treatment**	without supplements	1,52	695,92
		one feeding	1,52	695,92
		two feedings	1,67	916,98
	Rhizohumin	without supplements	1,73	1012,70
		one feeding	1,75	1110,50
		two feedings	2,04	1541,55
	Emistym C	without supplements	1,67	916,98
		one feeding	1,73	1012,70
		two feedings	2,11	1594,45
	Rhizohumin + Emistym C	without supplements	2,16	1491,36
		one feeding	2,22	1761,59
		two feedings	2,35	1829,73

Notes: * – Emistym C; ** - CONTROL.

The use of the biological preparation Rizogumin with the use of two foliar feedings of Emistym C contributed to an increase in the chlorophyll content in lupine of the Veresnevy white variety by 0.3 mg/g, and in the Makarivsky variety by 0.59 mg/g compared to the control. On both varieties of white lupine, pre-sowing treatment of seeds with a complex of drugs Rhizogumin and Emistym C followed by foliar feeding of plants with Emistym C provided the highest pigment content in the experiment, both in terms of raw weight and in terms of unit area. In plants of the Veresnevyi variety, these indicators were 2.87 mg/g and 4802.12 mg/m², in the Makarivskyi variety - 2.35 mg/g and 1829.73 mg/m². It was noted that the intensity of dry matter accumulation during the growing season of white lupine varieties depended on the studied factors, namely pre-sowing seed treatment and foliar fertilization. Observations of the dynamics of dry matter accumulation in white lupine varieties showed that the maximum yield is formed in the phase of physiological maturity (Table 3).

Table 3. Dynamics of accumulation of dry matter of white lupine varieties depending on biological preparations, g/plant (average for 2018-2020).

Factors			Air-dry mass of the aerial part of the plant, g/plant		
variety	pre-sowing seed processing	foliar feeding*	budding	full bloom	full ripeness
Veresnevyi	Without pre-sowing seed treatment**	without supplements**	3,13	4,34	8,61
		one feeding	3,15	4,45	8,64
		two feedings	3,17	4,56	8,67

	Rhizohumin	without supplements	3,16	4,51	8,70
		one feeding	3,18	4,62	8,74
		two feedings	3,20	4,73	8,83
	Emistym C	without supplements	3,19	4,84	8,71
		one feeding	3,21	4,95	8,76
		two feedings	3,23	5,06	8,81
	Rhizohumin + Emistym C	without supplements	3,25	5,07	8,80
		one feeding	3,28	5,41	8,84
		two feedings	3,31	5,84	8,99
Makarovskyi	Without pre-sowing seed treatment ^{**}	without supplements	2,84	3,86	8,26
		one feeding	2,86	3,95	8,30
		two feedings	2,88	4,04	8,34
	Rhizohumin	without supplements	2,87	4,13	8,40
		one feeding	2,89	4,22	8,46
		two feedings	2,91	4,31	8,50
	Emistym C	without supplements	2,90	4,25	8,32
		one feeding	2,92	4,34	8,37
		two feedings	2,94	4,43	8,41
	Rhizohumin + Emistym C	without supplements	2,96	4,52	8,50
		one feeding	2,99	4,61	8,53
		two feedings	3,02	4,70	8,61

Notes: * – Emistym C; ** - CONTROL.

Thus, in the variants where the biopreparations Ryzogumin and Emistym C were applied in combination with two foliar top dressings Emistym C, the amount of dry matter was 8.99 t/ha in the Veresnevy variety, and 8.61 t/ha in the Makarivsky variety, which was 0.28 and by 0.35 t/ha more than in the variants without application of pre-sowing treatment and without foliar fertilization. Productivity is a complex complex feature that manifests itself on the basis of the functional activity of various plant organs that make up their morphological and physiological structure. Each organ (root, stem, leaf, seed) is formed at a certain stage of ontogenesis. Their vital activities are limited to different time periods and are regulated by the genetic apparatus of organisms in complex interaction with environmental conditions (Roggatz et al., 1999; Soltani et al., 2001; Ahmadi et al., 2010). In an average of 3 years, the research results indicate a significant impact of the researched technological methods of growing on the yield (Table 4).

Table 4. Grain yield of white lupine varieties depending on biological preparations, t/ha (average for 2018-2020).

variety	Factors		Years			Average
	pre-sowing seed processing	foliar feeding*	2018	2019	2020	
Veresnevyi	Without pre-sowing seed treatment ^{**}	without supplements ^{**}	3,08	3,24	2,55	2,96
		one feeding	3,13	3,35	2,59	3,02
		two feedings	3,18	3,42	2,62	3,17
	Rhizohumin	without supplements	3,15	3,71	2,90	3,25
		one feeding	3,31	3,88	2,94	3,38
		two feedings	3,40	3,90	3,05	3,45
	Emistym C	without supplements	3,10	3,68	2,82	3,20
		one feeding	3,20	3,74	2,86	3,27
		two feedings	3,31	3,81	2,93	3,35
	Rhizohumin + Emistym C	without supplements	3,08	3,62	2,88	3,19
		one feeding	3,12	3,85	3,01	3,32
		two feedings	3,58	4,10	3,15	3,61
Makarovskyi	Without pre-sowing seed treatment ^{**}	without supplements	2,69	2,74	2,46	2,63
		one feeding	2,78	2,81	2,54	2,71
		two feedings	2,90	2,93	2,62	2,81
	Rhizohumin	without supplements	3,00	3,13	2,51	2,88
		one feeding	3,14	3,31	2,72	3,05
		two feedings	3,20	3,45	2,80	3,15
	Emistym C	without supplements	2,68	2,78	2,28	2,58
		one feeding	2,71	2,85	2,32	2,62
		two feedings	2,80	2,90	2,50	2,73

Rhizohumin + Emystim C	without supplements	3,11	3,24	2,38	2,91
	one feeding	3,22	3,40	2,41	3,01
	two feedings	3,34	3,65	2,70	3,23
LSD _{0,5} t/ha: A-0,07; B-0,10; C-0,08; AB-0,14; AC-0,12; BC-0,17; ABC-0,24 2018 p. HIP _{0,5} t/ha: A-0,04; B-0,05; C-0,04; AB-0,07; AC-0,06; BC-0,08; ABC-0,12 2019 p. HIP _{0,5} t/ha: A-0,05; B-0,06; C-0,06; AB-0,09; AC-0,08; BC-0,11; ABC-0,16 2020 p. HIP _{0,5} t/ha: A-0,04; B-0,06; C-0,05; AB-0,08; AC-0,07; BC-0,10; ABC-0,14					

Notes: * – Emistym C; ** – CONTROL.

The maximum value of grain yield of white lupine of the Veresnevy variety was obtained in the variants of the experiment with pre-sowing seed treatment with the inoculant Rhizogumin and the growth stimulator Emistym C in combination with two foliar fertilizing Emistym C. At the same time, the amount of grain yield was 3.61 t/ha, and exceeded the control variant by 0.65 t/ha, and in the percentage ratio, respectively – 18%.

CONCLUSIONS

Vinnitsia National Agrarian University develops the latest technologies for growing legumes, including white lupine, with elements of biologization and implements the results in production, within the framework of the implementation of the state theme, which is carried out at the expense of the state budget on the topic: «Development of methods for improving cultivation technology legumes using biofertilizers, bacterial preparations, foliar feeding and physiologically active substances» (state registration number 0120U102034).

During 2018-2020, on the basis of the research farm Agronomichne" of the Vinnitsia National Agrarian University, field studies were conducted on the effect of treating white lupine with biological preparations. Pre-sowing treatment of seeds with nodule bacteria and a growth stimulator in combination with foliar fertilization influenced the formation of the value of growth and development processes, as well as quantitative and qualitative indicators of white lupine. On the basis of the obtained research results, with the aim of growing high yields of white lupine grain at the level of 3.23-3.61 t/ha, agroforming recommends:

- sowing the intensive variety of white lupine Veresnevy;
- carry out pre-sowing seed treatment with the bacterial preparation Rhizogumin (600 g per hectare of seed rate) and the growth stimulator Emistym C (10 ml per 1 ton of seeds) in combination with two foliar fertilizing Emistym C in two periods: the first - in the budding phase, the second - in phase of the beginning of grain pouring (15 ml/ha).

REFERENCES

1. Vdovenko, S. et al. (2018). Symbiotic potential of snap beans (*Phaseolus vulgaris* L.) depending on biological products in agrocoenosis of the right-bank forest-steppe of Ukraine. *Ukrainian Journal of Ecology*, 8 (3), 270–274;
2. Langewisch T. et al. (2017). The development and use of a molecular model for soybean maturity groups. *Bmc Plant Biology*, 17, 13;
3. Pancy`reva G. V. (2016). Doslidzhennya sortovy`x resursiv lyupy`nu bilogo (*Lupinus albus* L.) v Ukrayini. *Vinny`cya*, 4, 88-93;
4. Kaletnik, G., & Lutkovska, S. (2020). Innovative Environmental Strategy for Sustainable Development. *European Journal of Sustainable Development*, 9(2), 89, <https://doi.org/10.14207/ejsd.2020.v9n2p89>;
5. Nychyporovych, A.A. et al. (1961). Photosynthetic activity of plants in sowings. Moscow: USSR Academy of Sciences;
6. Chynchyk, O.S. (2013). Influence of fertilization system and methods of basic tillage on the formation of pea plant structure. Feed and feed production: interdepartmental thematic collection of scientific papers, 77, 123–127;
7. Didur I. et al. (2020). Agroecological rationale of technological methods of growing legumes. The scientific heritage, 52. P. 3–14;

8. Didur I., Bakhmat M., Chynchyk O., Pantsyreva H., Telekalo N., Tkachuk O. (2020) Substantiation of agroecological factors on soybean agrophytocenoses by analysis of variance of the Right-Bank Forest-Steppe in Ukraine. *Ukrainian Journal of Ecology*. 10 (5), 54-61;
9. Didur I., Chynchyk O., Pantsyreva H., Olifirovych S., Olifirovych V., Tkachuk O. (2021) Effect of fertilizers for *Phaseolus vulgaris* L. productivity in Western Forest-Steppe of Ukraine. *Ukrainian Journal of Ecology*. 11 (1), 419-424;
10. Kantolic A. (2013). Seed number responses to extended photoperiod and shading during reproductive stages in indeterminate soybean. *European Journal of Agronomy*, 51: 91–100;
11. Mazur, V. et al. (2021). Ecological suitability peas (*Pisum sativum*) varieties to climate change in Ukraine. *Agraarteadus*, 32 (2), 276–283, <https://doi.org/10.15159/jas.21.26>;
12. Honcharuk, I. et al. (2020). Efficiency of growing legumes crops in Ukraine. In :Integration of traditional and innovation processes of development of modern science: collective monograph (3rd ed). (Ed. A. Jankovska). Baltija Publishing, Riga, Latvia, 42– 65. DOI: 10.30525/978-9934-26-021- 6-31;
13. Kaletnik, G.M. (2010). Biofuels. Food, Energy and Environmental Security of Ukraine: Monograph. K: High-tech Press, 516;
14. Kaletnik, G. et al. (2020). The Waste-Free Production Development for the Energy Autonomy Formation of Ukrainian Agricultural Enterprises. *Journal of Environmental Management and Tourism*. 11(3): 513–522;
15. Zhu J.H. et al. (2019). Loss of Function of the E1-Like-b Gene Associates With Early Flowering Under Long-Day Conditions in Soybean. *Frontiers in Plant Science*, 9, 13;
16. Mazur, V.A. et al. (2019). Influence of the assimilation apparatus and productivity of white lupine plants. *Agronomy Research*, 17, 206-219;
17. Mazur, V. et al. (2020). Agroecological prospects of using corn hybrids for biogas production. *Agronomy Research*, 18, 205-219;
18. Puyu, V. et al. (2021). Socialand-Ecological Aspects of Forage Production Reform in Ukraine in the Early 21st Century. *European Journal of Sustainable Development*, 10(1), 221–228;
19. Roggatz, U. et al. (1999). Effects of nitrogen deprivation on cell division and expansion in leaves of *Ricinus communis* L. *Plant, Cell and Environment*, 22(1) : 81–89. DOI:1046/j.1365-3040.1999.00383.x;
20. Soltani, A. et al. (2001). A simulation study of chickpea crop response to limited irrigation in semiarid environment. *Agricultural Water Management*, 95: 171–181;
21. Ahmadi, H. et al. (2010). Effect of different levels of nitrogen fertilizer on yield, nitrate, accumulation and several quantitative attributes of five Iranian spinach accessions. *American Eurasian Journal of Agricultural and Environmental Science*, 8(4), 468–473;
22. Bandura, V. et al. (2019). Research on sunflower seeds drying process in a monolayer tray vibration dryer based on infrared radiation. *INMATEN. Agricultural Engineering*, 57(1), 233–242;
23. Rai R. et al. (2017). Exogenous application of ethrel and gibberellic acid stimulates physiological growth of late planted sugarcane with short growth period in subtropical India. *Journal of Plant Growth Regulation*, 36(2), 472-486;
24. Bulgakov V. et al. (2014). Mathematical model of vibration digging up of root crops from soil. *Agronomy Research*, 12 (1), 41-58;
25. Mazur V., Didur I., Tkachuk O., Pantsyreva H., Ovcharuk V. (2021) Agroecological stability of cultivars of sparsely distributed legumes in the context of climate change. *Scientific Horizons*. 24 (1), 54-60;
26. Mazur V., Tkachuk O., Pantsyreva H., Demchuk O. (2021) Quality of pea seeds and agroecological condition of soil when using structured water. *Scientific Horizons*. 24 (7), 53-60;
27. Mazur V., Tkachuk O., Pantsyreva H., Kupchuk I., Mordvaniuk M., Chynchyk O. (2021) Ecological suitability peas (*Pisum sativum*) varieties to climate change in Ukraine. *Agraarteadus. Journal of Agricultural Science*. 2 (32), 276-283;
28. Didur, I. et al. (2020). Substantiation of agroecological factors on soybean agrophytocenoses by analysis of variance of the Right-Bank Forest-Steppe in Ukraine. *Ukrainian Journal of Ecology*, 10 (5), 54-61;
29. Mazur, V. et al. (2019). Influence of the Photosynthetic Productivity and Seed Productivity of White Lupine Plants. *Ukrainian Journal of Ecology*, 9(4), 665-670;